

Journal Articles

- [1] G. Aad (ATLAS), “Observation of WWW Production in pp Collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **129**, 061803 (2022).
- [2] G. Aad (ATLAS), “A search for an unexpected asymmetry in the production of $e+\mu^-$ and $e-\mu^+$ pairs in proton–proton collisions recorded by the ATLAS detector at $s=13$ TeV”, *Phys. Lett. B* **830**, 137106 (2022).
- [3] G. Aad (ATLAS), “AtlFast3: the next generation of fast simulation in ATLAS”, *Comput. Softw. Big Sci.* **6**, 7 (2022).
- [4] G. Aad (ATLAS), “Constraints on Higgs boson production with large transverse momentum using $H \rightarrow b\bar{b}$ decays in the ATLAS detector”, *Phys. Rev. D* **105**, 092003 (2022).
- [5] G. Aad (ATLAS), “Constraints on Higgs boson properties using $WW^*(\rightarrow e\nu\mu\nu)jj$ production in 36.1 fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **82**, 622 (2022).
- [6] G. Aad (ATLAS), “Determination of the parton distribution functions of the proton using diverse ATLAS data from pp collisions at $\sqrt{s} = 7, 8$ and 13 TeV”, *Eur. Phys. J. C* **82**, 438 (2022).
- [7] G. Aad (ATLAS), “Direct constraint on the Higgs-charm coupling from a search for Higgs boson decays into charm quarks with the ATLAS detector”, *Eur. Phys. J. C* **82**, 717 (2022).
- [8] G. Aad (ATLAS), “Emulating the impact of additional proton–proton interactions in the ATLAS simulation by presampling sets of inelastic Monte Carlo events”, *Comput. Softw. Big Sci.* **6**, 3 (2022).
- [9] G. Aad (ATLAS), “Measurement of Higgs boson decay into b -quarks in associated production with a top-quark pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **06**, 097 (2022).
- [10] G. Aad (ATLAS), “Measurement of the c -jet mistagging efficiency in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS detector”, *Eur. Phys. J. C* **82**, 95 (2022).
- [11] G. Aad (ATLAS), “Measurement of the energy asymmetry in $t\bar{t}j$ production at 13 TeV with the ATLAS experiment and interpretation in the SMEFT framework”, *Eur. Phys. J. C* **82**, 374 (2022).
- [12] G. Aad (ATLAS), “Measurement of the energy response of the ATLAS calorimeter to charged pions from $W^\pm \rightarrow \tau^\pm(\rightarrow \pi^\pm\nu_\tau)\nu_\tau$ events in Run 2 data”, *Eur. Phys. J. C* **82**, 223 (2022).
- [13] G. Aad (ATLAS), “Measurement of the nuclear modification factor for muons from charm and bottom hadrons in Pb+Pb collisions at 5.02 TeV with the ATLAS detector”, *Phys. Lett. B* **829**, 137077 (2022).
- [14] G. Aad (ATLAS), “Measurement of the polarisation of single top quarks and antiquarks produced in the t -channel at $\sqrt{s} = 13$ TeV and bounds on the tWb dipole operator from the ATLAS experiment”, *JHEP* **11**, 040 (2022).
- [15] G. Aad (ATLAS), “Measurements of azimuthal anisotropies of jet production in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Phys. Rev. C* **105**, 064903 (2022).
- [16] G. Aad (ATLAS), “Measurements of differential cross-sections in top-quark pair events with a high transverse momentum top quark and limits on beyond the Standard Model contributions to top-quark pair production with the ATLAS detector at $\sqrt{s} = 13$ TeV”, *JHEP* **06**, 063 (2022).
- [17] G. Aad (ATLAS), “Measurements of Higgs boson production cross-sections in the $H \rightarrow \tau^+\tau^-$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **08**, 175 (2022).
- [18] G. Aad (ATLAS), “Measurements of jet observables sensitive to b -quark fragmentation in $t\bar{t}$ events at the LHC with the ATLAS detector”, *Phys. Rev. D* **106**, 032008 (2022).
- [19] G. Aad (ATLAS), “Measurements of the Higgs boson inclusive and differential fiducial cross-sections in the diphoton decay channel with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **08**, 027 (2022).
- [20] G. Aad (ATLAS), “Modelling and computational improvements to the simulation of single vector-boson plus jet processes for the ATLAS experiment”, *JHEP* **08**, 089 (2022).
- [21] G. Aad (ATLAS), “Observation of electroweak production of two jets in association with an isolated photon and missing transverse momentum, and search for a Higgs boson decaying into invisible particles at 13 TeV with the ATLAS detector”, *Eur. Phys. J. C* **82**, 105 (2022).
- [22] G. Aad (ATLAS), “Operation and performance of the ATLAS semiconductor tracker in LHC Run 2”, *JINST* **17**, P01013 (2022).
- [23] G. Aad (ATLAS), “Performance of the ATLAS Level-1 topological trigger in Run 2”, *Eur. Phys. J. C* **82**, 7 (2022).

- [24] G. Aad (ATLAS), “Search for associated production of a Z boson with an invisibly decaying Higgs boson or dark matter candidates at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **829**, 137066 (2022).
- [25] G. Aad (ATLAS), “Search for events with a pair of displaced vertices from long-lived neutral particles decaying into hadronic jets in the ATLAS muon spectrometer in pp collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **106**, 032005 (2022).
- [26] G. Aad (ATLAS), “Search for exotic decays of the Higgs boson into $b\bar{b}$ and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **01**, 063 (2022).
- [27] G. Aad (ATLAS), “Search for flavour-changing neutral-current interactions of a top quark and a gluon in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **82**, 334 (2022).
- [28] G. Aad (ATLAS), “Search for heavy particles in the b -tagged dijet mass distribution with additional b -tagged jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, *Phys. Rev. D* **105**, 012001 (2022).
- [29] G. Aad (ATLAS), “Search for Higgs boson decays into a pair of pseudoscalar particles in the $bb\mu\mu$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **105**, 012006 (2022).
- [30] G. Aad (ATLAS), “Search for Higgs boson pair production in the two bottom quarks plus two photons final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **106**, 052001 (2022).
- [31] G. Aad (ATLAS), “Search for Higgs bosons decaying into new spin-0 or spin-1 particles in four-lepton final states with the ATLAS detector with 139 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV”, *JHEP* **03**, 041 (2022).
- [32] G. Aad (ATLAS), “Search for invisible Higgs-boson decays in events with vector-boson fusion signatures using 139 fb^{-1} of proton-proton data recorded by the ATLAS experiment”, *JHEP* **08**, 104 (2022).
- [33] G. Aad (ATLAS), “Search for lepton-flavor-violation in Z -boson decays with τ -leptons with the ATLAS detector”, *Phys. Rev. Lett.* **127**, 271801 (2022).
- [34] G. Aad (ATLAS), “Search for long-lived charginos based on a disappearing-track signature using 136 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **82**, 606 (2022).
- [35] G. Aad (ATLAS), “Search for neutral long-lived particles in pp collisions at $\sqrt{s} = 13$ TeV that decay into displaced hadronic jets in the ATLAS calorimeter”, *JHEP* **06**, 005 (2022).
- [36] G. Aad (ATLAS), “Search for new phenomena in three- or four-lepton events in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **824**, 136832 (2022).
- [37] G. Aad (ATLAS), “Search for resonant pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **105**, 092002 (2022).
- [38] G. Aad (ATLAS), “Search for single production of a vectorlike T quark decaying into a Higgs boson and top quark with fully hadronic final states using the ATLAS detector”, *Phys. Rev. D* **105**, 092012 (2022).
- [39] G. Aad (ATLAS), “Search for type-III seesaw heavy leptons in leptonic final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **82**, 988 (2022).
- [40] G. Aad (ATLAS), “Study of $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **08**, 087 (2022).
- [41] G. Aad (ATLAS), “The ATLAS inner detector trigger performance in pp collisions at 13 TeV during LHC Run 2”, *Eur. Phys. J. C* **82**, 206 (2022).
- [42] G. Aad (ATLAS), “Two-particle Bose–Einstein correlations in pp collisions at $\sqrt{s} = 13$ TeV measured with the ATLAS detector at the LHC”, *Eur. Phys. J. C* **82**, 608 (2022).
- [43] S. Bailey, “Data and Analysis Preservation, Recasting, and Reinterpretation”, (2022).
- [44] A. Belloni, “Report of the Topical Group on Electroweak Precision Physics and Constraining New Physics for Snowmass 2021”, (2022).
- [45] W. Bhimij, “Snowmass 2021 Computational Frontier CompF4 Topical Group Report: Storage and Processing Resource Access”, (2022).
- [46] K. M. Black, “Muon Collider Forum Report”, (2022).
- [47] K. Cranmer, “Publishing statistical models: Getting the most out of particle physics experiments”, *SciPost Phys.* **12**, 037 (2022).
- [48] A. M. Deiana, “Applications and Techniques for Fast Machine Learning in Science”, *Front. Big Data* **5**, 787421 (2022).
- [49] J. Duarte, “FAIR AI Models in High Energy Physics”, (2022).

- [50] A. Elabd, “Graph Neural Networks for Charged Particle Tracking on FPGAs”, *Front. Big Data* **5**, 828666 (2022).
- [51] E. A. Huerta, “FAIR for AI: An interdisciplinary, international, inclusive, and diverse community building perspective”, (2022).
- [52] A. Khot, **M. S. Neubauer**, A. Roy, “A Detailed Study of Interpretability of Deep Neural Network based Top Taggers”, (2022).
- [53] F. Maltoni, “TF07 Snowmass Report: Theory of Collider Phenomena”, (2022).
- [54] B. Nachman, “Jets and Jet Substructure at Future Colliders”, *Front. in Phys.* **10**, 897719 (2022).
- [55] **M. Neubauer**, M. Feickert, M. Katare, A. Roy, “Deep Learning for the Matrix Element Method”, *PoS ICHEP2022*, 246 (2022).
- [56] **M. S. Neubauer**, A. Roy, Z. Wang, “Making Digital Objects FAIR in High Energy Physics: An Implementation for Universal FeynRules Output (UFO) Models”, (2022).
- [57] A. Roy, **M. S. Neubauer**, “Interpretability of an Interaction Network for identifying $H \rightarrow b\bar{b}$ jets”, *PoS ICHEP2022*, 223 (2022).
- [58] P. Shanahan, “Snowmass 2021 Computational Frontier CompF03 Topical Group Report: Machine Learning”, (2022).
- [59] M. Aaboud (ATLAS), “Measurement of the relative B_c^\pm/B^\pm production cross section with the ATLAS detector at $\sqrt{s} = 8$ TeV”, *Phys. Rev. D* **104**, 012010 (2021).
- [60] G. Aad (ATLAS), “A search for the decays of stopped long-lived particles at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **07**, 173 (2021).
- [61] G. Aad (ATLAS), “A search for the dimuon decay of the Standard Model Higgs boson with the ATLAS detector”, *Phys. Lett. B* **812**, 135980 (2021).
- [62] G. Aad (ATLAS), “Configuration and performance of the ATLAS b -jet triggers in Run 2”, *Eur. Phys. J. C* **81**, 1087 (2021).
- [63] G. Aad (ATLAS), “Determination of the parton distribution functions of the proton from ATLAS measurements of differential W and Z boson production in association with jets”, *JHEP* **07**, 223 (2021).
- [64] G. Aad (ATLAS), “Differential cross-section measurements for the electroweak production of dijets in association with a Z boson in proton–proton collisions at ATLAS”, *Eur. Phys. J. C* **81**, 163 (2021).
- [65] G. Aad (ATLAS), “Evidence for Higgs boson decays to a low-mass dilepton system and a photon in pp collisions at $s=13$ TeV with the ATLAS detector”, *Phys. Lett. B* **819**, 136412 (2021).
- [66] G. Aad (ATLAS), “Exclusive dimuon production in ultraperipheral Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ATLAS”, *Phys. Rev. C* **104**, 024906 (2021).
- [67] G. Aad (ATLAS), “Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **81**, 689 (2021).
- [68] G. Aad (ATLAS), “Longitudinal Flow Decorrelations in Xe+Xe Collisions at $\sqrt{s_{NN}} = 5.44$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **126**, 122301 (2021).
- [69] G. Aad (ATLAS), “Measurement of b -quark fragmentation properties in jets using the decay $B^\pm \rightarrow J/\psi K^\pm$ in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **12**, 131 (2021).
- [70] G. Aad (ATLAS), “Measurement of hadronic event shapes in high- p_T multijet final states at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **01**, [Erratum: *JHEP* 12, 053 (2021)], 188 (2021).
- [71] G. Aad (ATLAS), “Measurement of light-by-light scattering and search for axion-like particles with 2.2 nb $^{-1}$ of Pb+Pb data with the ATLAS detector”, *JHEP* **03**, [Erratum: *JHEP* 11, 050 (2021)], 243 (2021).
- [72] G. Aad (ATLAS), “Measurement of single top-quark production in association with a W boson in the single-lepton channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **81**, 720 (2021).
- [73] G. Aad (ATLAS), “Measurement of the CP -violating phase ϕ_s in $B_s^0 \rightarrow J/\psi\phi$ decays in ATLAS at 13 TeV”, *Eur. Phys. J. C* **81**, 342 (2021).
- [74] G. Aad (ATLAS), “Measurement of the associated production of a Higgs boson decaying into b -quarks with a vector boson at high transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **816**, 136204 (2021).
- [75] G. Aad (ATLAS), “Measurement of the jet mass in high transverse momentum $Z(\rightarrow b\bar{b})\gamma$ production at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Lett. B* **812**, 135991 (2021).
- [76] G. Aad (ATLAS), “Measurement of the production cross section of pairs of isolated photons in pp collisions at 13 TeV with the ATLAS detector”, *JHEP* **11**, 169 (2021).
- [77] G. Aad (ATLAS), “Measurement of the $t\bar{t}t\bar{t}$ production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **11**, 118 (2021).

- [78] G. Aad (ATLAS), “Measurements of $W^+W^- + \geq 1$ jet production cross-sections in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **06**, 003 (2021).
- [79] G. Aad (ATLAS), “Measurements of WH and ZH production in the $H \rightarrow b\bar{b}$ decay channel in pp collisions at 13 TeV with the ATLAS detector”, Eur. Phys. J. C **81**, 178 (2021).
- [80] G. Aad (ATLAS), “Measurements of differential cross-sections in four-lepton events in 13 TeV proton-proton collisions with the ATLAS detector”, JHEP **07**, 005 (2021).
- [81] G. Aad (ATLAS), “Measurements of Higgs bosons decaying to bottom quarks from vector boson fusion production with the ATLAS experiment at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **81**, 537 (2021).
- [82] G. Aad (ATLAS), “Measurements of sensor radiation damage in the ATLAS inner detector using leakage currents”, JINST **16**, P08025 (2021).
- [83] G. Aad (ATLAS), “Measurements of the inclusive and differential production cross sections of a top-quark–antiquark pair in association with a Z boson at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **81**, 737 (2021).
- [84] G. Aad (ATLAS), “Measurements of top-quark pair single- and double-differential cross-sections in the all-hadronic channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, JHEP **01**, 033 (2021).
- [85] G. Aad (ATLAS), “Medium-Induced Modification of Z-Tagged Charged Particle Yields in $Pb + Pb$ Collisions at 5.02 TeV with the ATLAS Detector”, Phys. Rev. Lett. **126**, 072301 (2021).
- [86] G. Aad (ATLAS), “Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **81**, 578 (2021).
- [87] G. Aad (ATLAS), “Observation of photon-induced W^+W^- production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, Phys. Lett. B **816**, 136190 (2021).
- [88] G. Aad (ATLAS), “Optimisation of large-radius jet reconstruction for the ATLAS detector in 13 TeV proton–proton collisions”, Eur. Phys. J. C **81**, 334 (2021).
- [89] G. Aad (ATLAS), “Performance of the ATLAS RPC detector and Level-1 muon barrel trigger at $\sqrt{s} = 13$ TeV”, JINST **16**, P07029 (2021).
- [90] G. Aad (ATLAS), “Search for a heavy Higgs boson decaying into a Z boson and another heavy Higgs boson in the $\ell\ell b\bar{b}$ and $\ell\ell WW$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **81**, 396 (2021).
- [91] G. Aad (ATLAS), “Search for bottom-squark pair production in pp collision events at $\sqrt{s} = 13$ TeV with hadronically decaying τ -leptons, b -jets and missing transverse momentum using the ATLAS detector”, Phys. Rev. D **104**, 032014 (2021).
- [92] G. Aad (ATLAS), “Search for charged Higgs bosons decaying into a top quark and a bottom quark at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **06**, 145 (2021).
- [93] G. Aad (ATLAS), “Search for charged-lepton-flavour violation in Z-boson decays with the ATLAS detector”, Nature Phys. **17**, 819–825 (2021).
- [94] G. Aad (ATLAS), “Search for chargino–neutralino pair production in final states with three leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **81**, 1118 (2021).
- [95] G. Aad (ATLAS), “Search for charginos and neutralinos in final states with two boosted hadronically decaying bosons and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **104**, 112010 (2021).
- [96] G. Aad (ATLAS), “Search for dark matter in association with an energetic photon in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **02**, 226 (2021).
- [97] G. Aad (ATLAS), “Search for dark matter in events with missing transverse momentum and a Higgs boson decaying into two photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **10**, 013 (2021).
- [98] G. Aad (ATLAS), “Search for Dark Matter Produced in Association with a Dark Higgs Boson Decaying into $W^\pm W^\mp$ or ZZ in Fully Hadronic Final States from $\sqrt{s} = 13$ TeV pp Collisions Recorded with the ATLAS Detector”, Phys. Rev. Lett. **126**, 121802 (2021).
- [99] G. Aad (ATLAS), “Search for dark matter produced in association with a single top quark in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **81**, 860 (2021).
- [100] G. Aad (ATLAS), “Search for dark matter produced in association with a Standard Model Higgs boson decaying into b-quarks using the full Run 2 dataset from the ATLAS detector”, JHEP **11**, 209 (2021).
- [101] G. Aad (ATLAS), “Search for Displaced Leptons in $\sqrt{s} = 13$ TeV pp Collisions with the ATLAS Detector”, Phys. Rev. Lett. **127**, 051802 (2021).

- [102] G. Aad (ATLAS), “Search for doubly and singly charged Higgs bosons decaying into vector bosons in multi-lepton final states with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **06**, 146 (2021).
- [103] G. Aad (ATLAS), “Search for exotic decays of the Higgs boson into long-lived particles in pp collisions at $\sqrt{s} = 13$ TeV using displaced vertices in the ATLAS inner detector”, JHEP **11**, 229 (2021).
- [104] G. Aad (ATLAS), “Search for heavy resonances decaying into a pair of Z bosons in the $\ell^+\ell^-\ell'^+\ell'^-$ and $\ell^+\ell^-\nu\bar{\nu}$ final states using 139 fb^{-1} of proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **81**, 332 (2021).
- [105] G. Aad (ATLAS), “Search for Higgs boson production in association with a high-energy photon via vector-boson fusion with decay into bottom quark pairs at $\sqrt{s}=13$ TeV with the ATLAS detector”, JHEP **03**, 268 (2021).
- [106] G. Aad (ATLAS), “Search for new phenomena in pp collisions in final states with tau leptons, b-jets, and missing transverse momentum with the ATLAS detector”, Phys. Rev. D **104**, 112005 (2021).
- [107] G. Aad (ATLAS), “Search for new phenomena in events with an energetic jet and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **103**, 112006 (2021).
- [108] G. Aad (ATLAS), “Search for new phenomena in events with two opposite-charge leptons, jets and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **04**, 165 (2021).
- [109] G. Aad (ATLAS), “Search for new phenomena in final states with b -jets and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, JHEP **05**, 093 (2021).
- [110] G. Aad (ATLAS), “Search for New Phenomena in Final States with Two Leptons and One or No b -Tagged Jets at $\sqrt{s} = 13$ TeV Using the ATLAS Detector”, Phys. Rev. Lett. **127**, 141801 (2021).
- [111] G. Aad (ATLAS), “Search for new phenomena with top quark pairs in final states with one lepton, jets, and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **04**, 174 (2021).
- [112] G. Aad (ATLAS), “Search for pair production of scalar leptoquarks decaying into first- or second-generation leptons and top quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **81**, 313 (2021).
- [113] G. Aad (ATLAS), “Search for pair production of third-generation scalar leptoquarks decaying into a top quark and a τ -lepton in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **06**, 179 (2021).
- [114] G. Aad (ATLAS), “Search for phenomena beyond the Standard Model in events with large b -jet multiplicity using the ATLAS detector at the LHC”, Eur. Phys. J. C **81**, [Erratum: Eur.Phys.J.C 81, 249 (2021)], 11 (2021).
- [115] G. Aad (ATLAS), “Search for R-parity-violating supersymmetry in a final state containing leptons and many jets with the ATLAS experiment using $\sqrt{s} = 13\text{TeV}$ proton-proton collision data”, Eur. Phys. J. C **81**, 1023 (2021).
- [116] G. Aad (ATLAS), “Search for resonances decaying into photon pairs in 139 fb^{-1} of pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector”, Phys. Lett. B **822**, 136651 (2021).
- [117] G. Aad (ATLAS), “Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb^{-1} of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, JHEP **02**, 143 (2021).
- [118] G. Aad (ATLAS), “Search for squarks and gluinos in final states with one isolated lepton, jets, and missing transverse momentum at $\sqrt{s} = 13$ with the ATLAS detector”, Eur. Phys. J. C **81**, [Erratum: Eur.Phys.J.C 81, 956 (2021)], 600 (2021).
- [119] G. Aad (ATLAS), “Search for supersymmetry in events with four or more charged leptons in 139 fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, JHEP **07**, 167 (2021).
- [120] G. Aad (ATLAS), “Search for trilepton resonances from chargino and neutralino pair production in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Phys. Rev. D **103**, 112003 (2021).
- [121] G. Aad (ATLAS), “Search for type-III seesaw heavy leptons in dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **81**, 218 (2021).
- [122] G. Aad (ATLAS), “Test of the universality of τ and μ lepton couplings in W -boson decays with the ATLAS detector”, Nature Phys. **17**, 813–818 (2021).
- [123] G. Aad (ATLAS), “The ATLAS Fast TracKer system”, JINST **16**, P07006 (2021).

- [124] G. Aad (ATLAS), “Two-particle azimuthal correlations in photonuclear ultraperipheral Pb+Pb collisions at 5.02 TeV with ATLAS”, *Phys. Rev. C* **104**, 014903 (2021).
- [125] Y. Chen, “A FAIR and AI-ready Higgs boson decay dataset”, 10.1038/s41597-021-01109-0 (2021).
- [126] K. Choi, A. Eckart, B. Galewsky, R. Gardner, **M. S. Neubauer**, P. Onyisi, M. Proffitt, I. Vukotic, G. T. Watts, “Towards Real-World Applications of ServiceX, an Analysis Data Transformation System”, *EPJ Web Conf.* **251**, 02053 (2021).
- [127] G. DeZoort, S. Thais, J. Duarte, V. Razavimaleki, M. Atkinson, I. Ojalvo, **M. Neubauer**, P. Elmer, “Charged Particle Tracking via Edge-Classifying Interaction Networks”, *Comput. Softw. Big Sci.* **5**, 26 (2021).
- [128] X. Ju, “Performance of a geometric deep learning pipeline for HL-LHC particle tracking”, *Eur. Phys. J. C* **81**, 876 (2021).
- [129] M. Aaboud (ATLAS), “Determination of jet calibration and energy resolution in proton-proton collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **80**, 1104 (2020).
- [130] M. Aaboud (ATLAS), “Fluctuations of anisotropic flow in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *JHEP* **01**, 051 (2020).
- [131] M. Aaboud (ATLAS), “Measurement of J/ψ production in association with a W boson with pp data at 8 TeV”, *JHEP* **01**, 095 (2020).
- [132] M. Aaboud (ATLAS), “Measurement of long-range two-particle azimuthal correlations in Z-boson tagged pp collisions at $\sqrt{s}=8$ and 13 TeV”, *Eur. Phys. J. C* **80**, 64 (2020).
- [133] M. Aaboud (ATLAS), “Measurements of top-quark pair spin correlations in the $e\mu$ channel at $\sqrt{s} = 13$ TeV using pp collisions in the ATLAS detector”, *Eur. Phys. J. C* **80**, 754 (2020).
- [134] G. Aad (ATLAS), “CP Properties of Higgs Boson Interactions with Top Quarks in the $t\bar{t}H$ and tH Processes Using $H \rightarrow \gamma\gamma$ with the ATLAS Detector”, *Phys. Rev. Lett.* **125**, 061802 (2020).
- [135] G. Aad (ATLAS), “Z boson production in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured by the ATLAS experiment”, *Phys. Lett. B* **802**, 135262 (2020).
- [136] G. Aad (ATLAS), “A search for the $Z\gamma$ decay mode of the Higgs boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **809**, 135754 (2020).
- [137] G. Aad (ATLAS), “Alignment of the ATLAS Inner Detector in Run-2”, *Eur. Phys. J. C* **80**, 1194 (2020).
- [138] G. Aad (ATLAS), “ATLAS data quality operations and performance for 2015–2018 data-taking”, *JINST* **15**, P04003 (2020).
- [139] G. Aad (ATLAS), “Combination of searches for Higgs boson pairs in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **800**, 135103 (2020).
- [140] G. Aad (CMS, ATLAS), “Combination of the W boson polarization measurements in top quark decays using ATLAS and CMS data at $\sqrt{s} = 8$ TeV”, *JHEP* **08**, 051 (2020).
- [141] G. Aad (ATLAS), “Combined measurements of Higgs boson production and decay using up to 80 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment”, *Phys. Rev. D* **101**, 012002 (2020).
- [142] G. Aad (ATLAS), “Dijet resonance search with weak supervision using $\sqrt{s} = 13$ TeV pp collisions in the ATLAS detector”, *Phys. Rev. Lett.* **125**, 131801 (2020).
- [143] G. Aad (ATLAS), “Evidence for $t\bar{t}t\bar{t}$ production in the multilepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80**, 1085 (2020).
- [144] G. Aad (ATLAS), “Evidence for electroweak production of two jets in association with a $Z\gamma$ pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **803**, 135341 (2020).
- [145] G. Aad (ATLAS), “Higgs boson production cross-section measurements and their EFT interpretation in the 4ℓ decay channel at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80**, [Erratum: *Eur.Phys.J.C* 81, 29 (2021), Erratum: *Eur.Phys.J.C* 81, 398 (2021)], 957 (2020).
- [146] G. Aad (ATLAS), “Measurement of azimuthal anisotropy of muons from charm and bottom hadrons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **124**, 082301 (2020).
- [147] G. Aad (ATLAS), “Measurement of azimuthal anisotropy of muons from charm and bottom hadrons in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Phys. Lett. B* **807**, 135595 (2020).
- [148] G. Aad (ATLAS), “Measurement of differential cross sections for single diffractive dissociation in $\sqrt{s} = 8$ TeV pp collisions using the ATLAS ALFA spectrometer”, *JHEP* **02**, [Erratum: *JHEP* 10, 182 (2020)], 042 (2020).
- [149] G. Aad (ATLAS), “Measurement of isolated-photon plus two-jet production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **03**, 179 (2020).

- [150] G. Aad (ATLAS), “Measurement of soft-drop jet observables in pp collisions with the ATLAS detector at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **101**, 052007 (2020).
- [151] G. Aad (ATLAS), “Measurement of the $t\bar{t}$ production cross-section and lepton differential distributions in $e\mu$ dilepton events from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80**, 528 (2020).
- [152] G. Aad (ATLAS), “Measurement of the $t\bar{t}$ production cross-section in the lepton+jets channel at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, *Phys. Lett. B* **810**, 135797 (2020).
- [153] G. Aad (ATLAS), “Measurement of the $Z(\rightarrow \ell^+\ell^-)\gamma$ production cross-section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **03**, 054 (2020).
- [154] G. Aad (ATLAS), “Measurement of the azimuthal anisotropy of charged-particle production in $Xe + Xe$ collisions at $\sqrt{s_{NN}} = 5.44$ TeV with the ATLAS detector”, *Phys. Rev. C* **101**, 024906 (2020).
- [155] G. Aad (ATLAS), “Measurement of the Lund Jet Plane Using Charged Particles in 13 TeV Proton-Proton Collisions with the ATLAS Detector”, *Phys. Rev. Lett.* **124**, 222002 (2020).
- [156] G. Aad (ATLAS), “Measurement of the transverse momentum distribution of Drell–Yan lepton pairs in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80**, 616 (2020).
- [157] G. Aad (ATLAS), “Measurements of inclusive and differential cross-sections of combined $t\bar{t}\gamma$ and $tW\gamma$ production in the $e\mu$ channel at 13 TeV with the ATLAS detector”, *JHEP* **09**, 049 (2020).
- [158] G. Aad (ATLAS), “Measurements of the Higgs boson inclusive and differential fiducial cross sections in the 4ℓ decay channel at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **80**, 942 (2020).
- [159] G. Aad (ATLAS), “Measurements of the production cross-section for a Z boson in association with b -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **07**, 044 (2020).
- [160] G. Aad (ATLAS), “Observation and Measurement of Forward Proton Scattering in Association with Lepton Pairs Produced via the Photon Fusion Mechanism at ATLAS”, *Phys. Rev. Lett.* **125**, 261801 (2020).
- [161] G. Aad (ATLAS), “Observation of electroweak production of two jets and a Z -boson pair”, (2020).
- [162] G. Aad (ATLAS), “Observation of the associated production of a top quark and a Z boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **07**, 124 (2020).
- [163] G. Aad (ATLAS), “Operation of the ATLAS trigger system in Run 2”, *JINST* **15**, P10004 (2020).
- [164] G. Aad (ATLAS), “Performance of electron and photon triggers in ATLAS during LHC Run 2”, *Eur. Phys. J. C* **80**, 47 (2020).
- [165] G. Aad (ATLAS), “Performance of the ATLAS muon triggers in Run 2”, *JINST* **15**, P09015 (2020).
- [166] G. Aad (ATLAS), “Performance of the missing transverse momentum triggers for the ATLAS detector during Run-2 data taking”, *JHEP* **08**, 080 (2020).
- [167] G. Aad (ATLAS), “Performance of the upgraded PreProcessor of the ATLAS Level-1 Calorimeter Trigger”, *JINST* **15**, P11016 (2020).
- [168] G. Aad (ATLAS), “Reconstruction and identification of boosted di- τ systems in a search for Higgs boson pairs using 13 TeV proton-proton collision data in ATLAS”, *JHEP* **11**, 163 (2020).
- [169] G. Aad (ATLAS), “Search for $t\bar{t}$ resonances in fully hadronic final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **10**, 061 (2020).
- [170] G. Aad (ATLAS), “Search for a scalar partner of the top quark in the all-hadronic $t\bar{t}$ plus missing transverse momentum final state at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80**, 737 (2020).
- [171] G. Aad (ATLAS), “Search for chargino-neutralino production with mass splittings near the electroweak scale in three-lepton final states in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **101**, 072001 (2020).
- [172] G. Aad (ATLAS), “Search for dijet resonances in events with an isolated charged lepton using $\sqrt{s} = 13$ TeV proton-proton collision data collected by the ATLAS detector”, *JHEP* **06**, 151 (2020).
- [173] G. Aad (ATLAS), “Search for direct production of electroweakinos in final states with missing transverse momentum and a Higgs boson decaying into photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **10**, 005 (2020).
- [174] G. Aad (ATLAS), “Search for direct production of electroweakinos in final states with one lepton, missing transverse momentum and a Higgs boson decaying into two b -jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80**, 691 (2020).
- [175] G. Aad (ATLAS), “Search for direct stau production in events with two hadronic τ -leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **101**, 032009 (2020).

- [176] G. Aad (ATLAS), “Search for displaced vertices of oppositely charged leptons from decays of long-lived particles in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **801**, 135114 (2020).
- [177] G. Aad (ATLAS), “Search for electroweak production of charginos and sleptons decaying into final states with two leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector”, Eur. Phys. J. C **80**, 123 (2020).
- [178] G. Aad (ATLAS), “Search for flavour-changing neutral currents in processes with one top quark and a photon using 81 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, Phys. Lett. B **800**, 135082 (2020).
- [179] G. Aad (ATLAS), “Search for heavy diboson resonances in semileptonic final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **80**, 1165 (2020).
- [180] G. Aad (ATLAS), “Search for heavy Higgs bosons decaying into two tau leptons with the ATLAS detector using pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. Lett. **125**, 051801 (2020).
- [181] G. Aad (ATLAS), “Search for heavy neutral Higgs bosons produced in association with b -quarks and decaying into b -quarks at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **102**, 032004 (2020).
- [182] G. Aad (ATLAS), “Search for heavy resonances decaying into a photon and a hadronically decaying Higgs boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. Lett. **125**, 251802 (2020).
- [183] G. Aad (ATLAS), “Search for Higgs Boson Decays into a Z Boson and a Light Hadronically Decaying Resonance Using 13 TeV pp Collision Data from the ATLAS Detector”, Phys. Rev. Lett. **125**, 221802 (2020).
- [184] G. Aad (ATLAS), “Search for Higgs boson decays into two new low-mass spin-0 particles in the $4b$ channel with the ATLAS detector using pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **102**, 112006 (2020).
- [185] G. Aad (ATLAS), “Search for light long-lived neutral particles produced in pp collisions at $\sqrt{s} = 13$ TeV and decaying into collimated leptons or light hadrons with the ATLAS detector”, Eur. Phys. J. C **80**, 450 (2020).
- [186] G. Aad (ATLAS), “Search for long-lived neutral particles produced in pp collisions at $\sqrt{s} = 13$ TeV decaying into displaced hadronic jets in the ATLAS inner detector and muon spectrometer”, Phys. Rev. D **101**, 052013 (2020).
- [187] G. Aad (ATLAS), “Search for long-lived, massive particles in events with a displaced vertex and a muon with large impact parameter in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **102**, 032006 (2020).
- [188] G. Aad (ATLAS), “Search for Magnetic Monopoles and Stable High-Electric-Charge Objects in 13 TeV Proton-Proton Collisions with the ATLAS Detector”, Phys. Rev. Lett. **124**, 031802 (2020).
- [189] G. Aad (ATLAS), “Search for new non-resonant phenomena in high-mass dilepton final states with the ATLAS detector”, JHEP **11**, [Erratum: JHEP 04, 142 (2021)], 005 (2020).
- [190] G. Aad (ATLAS), “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s} = 13$ TeV proton-proton collisions recorded by ATLAS in Run 2 of the LHC”, JHEP **10**, 062 (2020).
- [191] G. Aad (ATLAS), “Search for new resonances in mass distributions of jet pairs using 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **03**, 145 (2020).
- [192] G. Aad (ATLAS), “Search for non-resonant Higgs boson pair production in the $b\bar{b}\nu\ell\nu$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **801**, 135145 (2020).
- [193] G. Aad (ATLAS), “Search for pairs of scalar leptoquarks decaying into quarks and electrons or muons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, JHEP **10**, 112 (2020).
- [194] G. Aad (ATLAS), “Search for resonances decaying into a weak vector boson and a Higgs boson in the fully hadronic final state produced in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **102**, 112008 (2020).
- [195] G. Aad (ATLAS), “Search for squarks and gluinos in final states with same-sign leptons and jets using 139 fb^{-1} of data collected with the ATLAS detector”, JHEP **06**, 046 (2020).
- [196] G. Aad (ATLAS), “Search for the $HH \rightarrow b\bar{b}b\bar{b}$ process via vector-boson fusion production using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **07**, [Erratum: JHEP 01, 145 (2021), Erratum: JHEP 05, 207 (2021)], 108 (2020).
- [197] G. Aad (ATLAS), “Search for the Higgs boson decays $H \rightarrow ee$ and $H \rightarrow e\mu$ in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **801**, 135148 (2020).

- [198] G. Aad (ATLAS), “Search for top squarks in events with a Higgs or Z boson using 139 fb^{-1} of pp collision data at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, *Eur. Phys. J. C* **80**, 1080 (2020).
- [199] G. Aad (ATLAS), “Searches for electroweak production of supersymmetric particles with compressed mass spectra in $\sqrt{s} = 13 \text{ TeV}$ pp collisions with the ATLAS detector”, *Phys. Rev. D* **101**, 052005 (2020).
- [200] G. Aad (ATLAS), “Searches for lepton-flavour-violating decays of the Higgs boson in $\sqrt{s} = 13 \text{ TeV}$ pp collisions with the ATLAS detector”, *Phys. Lett. B* **800**, 135069 (2020).
- [201] G. Aad (ATLAS), “Test of CP invariance in vector-boson fusion production of the Higgs boson in the $H \rightarrow \tau\tau$ channel in proton–proton collisions at $s=13\text{TeV}$ with the ATLAS detector”, *Phys. Lett. B* **805**, 135426 (2020).
- [202] G. Aad (ATLAS), “Transverse momentum and process dependent azimuthal anisotropies in $\sqrt{s_{NN}} = 8.16 \text{ TeV}$ $p+\text{Pb}$ collisions with the ATLAS detector”, *Eur. Phys. J. C* **80**, 73 (2020).
- [203] J. Alison, “Higgs boson potential at colliders: Status and perspectives”, *Rev. Phys.* **5**, edited by B. Di Micco, M. Gouzevitch, J. Mazzitelli, and C. Vernieri, 100045 (2020).
- [204] B. Galewsky, R. Gardner, L. Gray, **M. Neubauer**, J. Pivarski, M. Proffitt, I. Vukotic, G. Watts, M. Weinberg, “ServiceX A Distributed, Caching, Columnar Data Delivery Service”, *EPJ Web Conf.* **245**, edited by C. Doglioni, D. Kim, G. A. Stewart, L. Silvestris, P. Jackson, and W. Kamleh, 04043 (2020).
- [205] R. Gardner, L. Bryant, **M. Neubauer**, F. Wuerthwein, J. Stephen, A. Chien, “The Scalable Systems Laboratory: a Platform for Software Innovation for HEP”, *EPJ Web Conf.* **245**, edited by C. Doglioni, D. Kim, G. A. Stewart, L. Silvestris, P. Jackson, and W. Kamleh, 05019 (2020).
- [206] M. Aaboud (ATLAS), “Search for scalar resonances decaying into $\mu^+\mu^-$ in events with and without b -tagged jets produced in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, *JHEP* **07**, 117 (2019).
- [207] M. Aaboud (ATLAS), “A search for pairs of highly collimated photon-jets in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, *Phys. Rev. D* **99**, 012008 (2019).
- [208] M. Aaboud (ATLAS), “A strategy for a general search for new phenomena using data-derived signal regions and its application within the ATLAS experiment”, *Eur. Phys. J. C* **79**, 120 (2019).
- [209] M. Aaboud (ATLAS), “Combination of searches for invisible Higgs boson decays with the ATLAS experiment”, *Phys. Rev. Lett.* **122**, 231801 (2019).
- [210] M. Aaboud (ATLAS, CMS), “Combinations of single-top-quark production cross-section measurements and $|f_{LV}V_{tb}|$ determinations at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS experiments”, *JHEP* **05**, 088 (2019).
- [211] M. Aaboud (ATLAS), “Comparison of Fragmentation Functions for Jets Dominated by Light Quarks and Gluons from pp and $\text{Pb}+\text{Pb}$ Collisions in ATLAS”, *Phys. Rev. Lett.* **123**, 042001 (2019).
- [212] M. Aaboud (ATLAS), “Constraints on mediator-based dark matter and scalar dark energy models using $\sqrt{s} = 13 \text{ TeV}$ pp collision data collected by the ATLAS detector”, *JHEP* **05**, 142 (2019).
- [213] M. Aaboud (ATLAS), “Correlated long-range mixed-harmonic fluctuations measured in pp , $p+\text{Pb}$ and low-multiplicity $\text{Pb}+\text{Pb}$ collisions with the ATLAS detector”, *Phys. Lett. B* **789**, 444–471 (2019).
- [214] M. Aaboud (ATLAS), “Cross-section measurements of the Higgs boson decaying into a pair of τ -leptons in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, *Phys. Rev. D* **99**, 072001 (2019).
- [215] M. Aaboud (ATLAS), “Dijet azimuthal correlations and conditional yields in pp and $p+\text{Pb}$ collisions at $s_{NN}=5.02\text{TeV}$ with the ATLAS detector”, *Phys. Rev. C* **100**, 034903 (2019).
- [216] M. Aaboud (ATLAS), “Electron and photon energy calibration with the ATLAS detector using 2015–2016 LHC proton-proton collision data”, *JINST* **14**, P03017 (2019).
- [217] M. Aaboud (ATLAS), “Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton-proton collision data at $\sqrt{s} = 13 \text{ TeV}$ ”, *Eur. Phys. J. C* **79**, 639 (2019).
- [218] M. Aaboud (ATLAS), “In situ calibration of large-radius jet energy and mass in 13 TeV proton–proton collisions with the ATLAS detector”, *Eur. Phys. J. C* **79**, 135 (2019).
- [219] M. Aaboud (ATLAS), “Measurement of $W^\pm Z$ production cross sections and gauge boson polarisation in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, *Eur. Phys. J. C* **79**, 535 (2019).
- [220] M. Aaboud (ATLAS), “Measurement of ZZ production in the $\ell\nu\nu$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ ”, *JHEP* **10**, 127 (2019).
- [221] M. Aaboud (ATLAS), “Measurement of fiducial and differential W^+W^- production cross-sections at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, *Eur. Phys. J. C* **79**, 884 (2019).

- [222] M. Aaboud (ATLAS), “Measurement of jet-substructure observables in top quark, W boson and light jet production in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **08**, 033 (2019).
- [223] M. Aaboud (ATLAS), “Measurement of photon–jet transverse momentum correlations in 5.02 TeV Pb + Pb and pp collisions with ATLAS”, Phys. Lett. B **789**, 167–190 (2019).
- [224] M. Aaboud (ATLAS), “Measurement of prompt photon production in $\sqrt{s_{NN}} = 8.16$ TeV p +Pb collisions with ATLAS”, Phys. Lett. B **796**, 230–252 (2019).
- [225] M. Aaboud (ATLAS), “Measurement of the $t\bar{t}Z$ and $t\bar{t}W$ cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **99**, 072009 (2019).
- [226] M. Aaboud (ATLAS), “Measurement of the four-lepton invariant mass spectrum in 13 TeV proton-proton collisions with the ATLAS detector”, JHEP **04**, 048 (2019).
- [227] M. Aaboud (ATLAS), “Measurement of the nuclear modification factor for inclusive jets in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, Phys. Lett. B **790**, 108–128 (2019).
- [228] M. Aaboud (ATLAS), “Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run 2 data collected in 2015 and 2016”, Eur. Phys. J. C **79**, 205 (2019).
- [229] M. Aaboud (ATLAS), “Measurement of the ratio of cross sections for inclusive isolated-photon production in pp collisions at $\sqrt{s} = 13$ and 8 TeV with the ATLAS detector”, JHEP **04**, 093 (2019).
- [230] M. Aaboud (ATLAS), “Measurement of the top quark mass in the $t\bar{t} \rightarrow$ lepton+jets channel from $\sqrt{s} = 8$ TeV ATLAS data and combination with previous results”, Eur. Phys. J. C **79**, 290 (2019).
- [231] M. Aaboud (ATLAS), “Measurement of $VH, H \rightarrow b\bar{b}$ production as a function of the vector-boson transverse momentum in 13 TeV pp collisions with the ATLAS detector”, JHEP **05**, 141 (2019).
- [232] M. Aaboud (ATLAS), “Measurements of W and Z boson production in pp collisions at $\sqrt{s} = 5.02$ TeV with the ATLAS detector”, Eur. Phys. J. C **79**, [Erratum: Eur.Phys.J.C 79, 374 (2019)], 128 (2019).
- [233] M. Aaboud (ATLAS), “Measurements of gluon-gluon fusion and vector-boson fusion Higgs boson production cross-sections in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **789**, 508–529 (2019).
- [234] M. Aaboud (ATLAS), “Measurements of inclusive and differential fiducial cross-sections of $t\bar{t}$ production with additional heavy-flavour jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **04**, 046 (2019).
- [235] M. Aaboud (ATLAS), “Measurements of inclusive and differential fiducial cross-sections of $t\bar{t}\gamma$ production in leptonic final states at $\sqrt{s} = 13$ TeV in ATLAS”, Eur. Phys. J. C **79**, 382 (2019).
- [236] M. Aaboud (ATLAS), “Modelling radiation damage to pixel sensors in the ATLAS detector”, JINST **14**, P06012 (2019).
- [237] M. Aaboud (ATLAS), “Observation of electroweak $W^\pm Z$ boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **793**, 469–492 (2019).
- [238] M. Aaboud (ATLAS), “Observation of electroweak production of a same-sign W boson pair in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. Lett. **123**, 161801 (2019).
- [239] M. Aaboud (ATLAS), “Performance of top-quark and W -boson tagging with ATLAS in Run 2 of the LHC”, Eur. Phys. J. C **79**, 375 (2019).
- [240] M. Aaboud (ATLAS), “Properties of $g \rightarrow b\bar{b}$ at small opening angles in pp collisions with the ATLAS detector at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **99**, 052004 (2019).
- [241] M. Aaboud (ATLAS), “Search for a right-handed gauge boson decaying into a high-momentum heavy neutrino and a charged lepton in pp collisions with the ATLAS detector at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **798**, 134942 (2019).
- [242] M. Aaboud (ATLAS), “Search for chargino and neutralino production in final states with a Higgs boson and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **100**, 012006 (2019).
- [243] M. Aaboud (ATLAS), “Search for doubly charged scalar bosons decaying into same-sign W boson pairs with the ATLAS detector”, Eur. Phys. J. C **79**, 58 (2019).
- [244] M. Aaboud (ATLAS), “Search for excited electrons singly produced in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment at the LHC”, Eur. Phys. J. C **79**, 803 (2019).
- [245] M. Aaboud (ATLAS), “Search for four-top-quark production in the single-lepton and opposite-sign dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **99**, 052009 (2019).

- [246] M. Aaboud (ATLAS), “Search for heavy charged long-lived particles in proton-proton collisions at $\sqrt{s} = 13$ TeV using an ionisation measurement with the ATLAS detector”, *Phys. Lett. B* **788**, 96–116 (2019).
- [247] M. Aaboud (ATLAS), “Search for heavy charged long-lived particles in the ATLAS detector in 36.1 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **99**, 092007 (2019).
- [248] M. Aaboud (ATLAS), “Search for heavy long-lived multicharged particles in proton-proton collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Rev. D* **99**, 052003 (2019).
- [249] M. Aaboud (ATLAS), “Search for heavy Majorana or Dirac neutrinos and right-handed W gauge bosons in final states with two charged leptons and two jets at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **01**, 016 (2019).
- [250] M. Aaboud (ATLAS), “Search for heavy particles decaying into a top-quark pair in the fully hadronic final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **99**, 092004 (2019).
- [251] M. Aaboud (ATLAS), “Search for Higgs boson decays into a pair of light bosons in the $b\bar{b}\mu\mu$ final state in pp collision at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **790**, 1–21 (2019).
- [252] M. Aaboud (ATLAS), “Search for Higgs boson pair production in the $b\bar{b}WW^*$ decay mode at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **04**, 092 (2019).
- [253] M. Aaboud (ATLAS), “Search for Higgs boson pair production in the $WW^{(*)}WW^{(*)}$ decay channel using ATLAS data recorded at $\sqrt{s} = 13$ TeV”, *JHEP* **05**, 124 (2019).
- [254] M. Aaboud (ATLAS), “Search for invisible Higgs boson decays in vector boson fusion at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **793**, 499–519 (2019).
- [255] M. Aaboud (ATLAS), “Search for large missing transverse momentum in association with one top-quark in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **05**, 041 (2019).
- [256] M. Aaboud (ATLAS), “Search for light resonances decaying to boosted quark pairs and produced in association with a photon or a jet in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **788**, 316–335 (2019).
- [257] M. Aaboud (ATLAS), “Search for long-lived neutral particles in pp collisions at $\sqrt{s} = 13$ TeV that decay into displaced hadronic jets in the ATLAS calorimeter”, *Eur. Phys. J. C* **79**, 481 (2019).
- [258] M. Aaboud (ATLAS), “Search for long-lived particles in final states with displaced dimuon vertices in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **99**, 012001 (2019).
- [259] M. Aaboud (ATLAS), “Search for long-lived particles produced in pp collisions at $\sqrt{s} = 13$ TeV that decay into displaced hadronic jets in the ATLAS muon spectrometer”, *Phys. Rev. D* **99**, 052005 (2019).
- [260] M. Aaboud (ATLAS), “Search for low-mass resonances decaying into two jets and produced in association with a photon using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **795**, 56–75 (2019).
- [261] M. Aaboud (ATLAS), “Search for pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **01**, 030 (2019).
- [262] M. Aaboud (ATLAS), “Search for single production of vector-like quarks decaying into Wb in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **05**, 164 (2019).
- [263] M. Aaboud (ATLAS), “Search for squarks and gluinos in final states with hadronically decaying τ -leptons, jets, and missing transverse momentum using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **99**, 012009 (2019).
- [264] M. Aaboud (ATLAS), “Search for the Production of a Long-Lived Neutral Particle Decaying within the ATLAS Hadronic Calorimeter in Association with a Z Boson from pp Collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **122**, 151801 (2019).
- [265] M. Aaboud (ATLAS), “Search for top-quark decays $t \rightarrow Hq$ with 36 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **05**, 123 (2019).
- [266] M. Aaboud (ATLAS), “Search for vector-boson resonances decaying to a top quark and bottom quark in the lepton plus jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **788**, 347–370 (2019).
- [267] M. Aaboud (ATLAS), “Searches for scalar leptoquarks and differential cross-section measurements in dilepton-dijet events in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV with the ATLAS experiment”, *Eur. Phys. J. C* **79**, 733 (2019).
- [268] M. Aaboud (ATLAS), “Searches for third-generation scalar leptoquarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *JHEP* **06**, 144 (2019).

- [269] M. Aaboud (ATLAS), “Study of the hard double-parton scattering contribution to inclusive four-lepton production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **790**, 595–614 (2019).
- [270] M. Aaboud (ATLAS), “Study of the rare decays of B_s^0 and B^0 mesons into muon pairs using data collected during 2015 and 2016 with the ATLAS detector”, *JHEP* **04**, 098 (2019).
- [271] G. Aad (ATLAS), “ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **79**, 970 (2019).
- [272] G. Aad (ATLAS), “Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data”, *JINST* **14**, P12006 (2019).
- [273] G. Aad (ATLAS), “Evidence for the production of three massive vector bosons with the ATLAS detector”, *Phys. Lett. B* **798**, 134913 (2019).
- [274] G. Aad (ATLAS), “Evidence for the production of three massive vectorbosons in pp collisions with the ATLAS detector”, *PoS DIS2019*, 135 (2019).
- [275] G. Aad (ATLAS), “Identification of boosted Higgs bosons decaying into b -quark pairs with the ATLAS detector at 13 TeV”, *Eur. Phys. J. C* **79**, 836 (2019).
- [276] G. Aad (ATLAS), “Measurement of K_S^0 and Λ^0 production in $t\bar{t}$ dileptonic events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **79**, 1017 (2019).
- [277] G. Aad (ATLAS), “Measurement of W^\pm boson production in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **79**, 935 (2019).
- [278] G. Aad (ATLAS), “Measurement of W^\pm -boson and Z-boson production cross-sections in pp collisions at $\sqrt{s} = 2.76$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **79**, 901 (2019).
- [279] G. Aad (ATLAS), “Measurement of angular and momentum distributions of charged particles within and around jets in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Phys. Rev. C* **100**, [Erratum: *Phys.Rev.C* 101, 059903 (2020)], 064901 (2019).
- [280] G. Aad (ATLAS), “Measurement of distributions sensitive to the underlying event in inclusive Z-boson production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **79**, 666 (2019).
- [281] G. Aad (ATLAS), “Measurement of flow harmonics correlations with mean transverse momentum in lead-lead and proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **79**, 985 (2019).
- [282] G. Aad (ATLAS), “Measurement of the cross-section and charge asymmetry of W bosons produced in proton–proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **79**, 760 (2019).
- [283] G. Aad (ATLAS), “Measurement of the inclusive cross-section for the production of jets in association with a Z boson in proton-proton collisions at 8 TeV using the ATLAS detector”, *Eur. Phys. J. C* **79**, 847 (2019).
- [284] G. Aad (ATLAS), “Measurement of the inclusive isolated-photon cross section in pp collisions at $\sqrt{s} = 13$ TeV using 36 fb^{-1} of ATLAS data”, *JHEP* **10**, 203 (2019).
- [285] G. Aad (ATLAS), “Measurement of the production cross section for a Higgs boson in association with a vector boson in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **798**, 134949 (2019).
- [286] G. Aad (ATLAS), “Measurement of the top-quark mass in $t\bar{t} + 1$ -jet events collected with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **11**, 150 (2019).
- [287] G. Aad (ATLAS), “Measurements of top-quark pair differential and double-differential cross-sections in the ℓ +jets channel with pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **79**, [Erratum: *Eur.Phys.J.C* 80, 1092 (2020)], 1028 (2019).
- [288] G. Aad (ATLAS), “Observation of light-by-light scattering in ultraperipheral Pb+Pb collisions with the ATLAS detector”, *Phys. Rev. Lett.* **123**, 052001 (2019).
- [289] G. Aad (ATLAS), “Properties of jet fragmentation using charged particles measured with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **100**, 052011 (2019).
- [290] G. Aad (ATLAS), “Resolution of the ATLAS muon spectrometer monitored drift tubes in LHC Run 2”, *JINST* **14**, P09011 (2019).
- [291] G. Aad (ATLAS), “Search for a heavy charged boson in events with a charged lepton and missing transverse momentum from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **100**, 052013 (2019).
- [292] G. Aad (ATLAS), “Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons, b -jets and missing transverse momentum”, *JHEP* **12**, 060 (2019).

- [293] G. Aad (ATLAS), “Search for diboson resonances in hadronic final states in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, JHEP **09**, [Erratum: JHEP 06, 042 (2020)], 091 (2019).
- [294] G. Aad (ATLAS), “Search for heavy neutral leptons in decays of W bosons produced in 13 TeV pp collisions using prompt and displaced signatures with the ATLAS detector”, JHEP **10**, 265 (2019).
- [295] G. Aad (ATLAS), “Search for high-mass dilepton resonances using 139 fb^{-1} of pp collision data collected at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Phys. Lett. B **796**, 68–87 (2019).
- [296] G. Aad (ATLAS), “Search for the electroweak diboson production in association with a high-mass dijet system in semileptonic final states in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Phys. Rev. D **100**, 032007 (2019).
- [297] J. Albrecht (HEP Software Foundation), “A Roadmap for HEP Software and Computing R&D for the 2020s”, Comput. Softw. Big Sci. **3**, 7 (2019).
- [298] E. A. Huerta, “Enabling real-time multi-messenger astrophysics discoveries with deep learning”, Nature Rev. Phys. **1**, 600–608 (2019).
- [299] E. A. Huerta, R. Haas, S. Jha, **M. Neubauer**, D. S. Katz, “Supporting High-Performance and High-Throughput Computing for Experimental Science”, Comput. Softw. Big Sci. **3**, 5 (2019).
- [300] M. Aaboud (ATLAS), “Combined measurement of differential and total cross sections in the $H \rightarrow \gamma\gamma$ and the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channels at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Phys. Lett. B **786**, 114–133 (2018).
- [301] M. Aaboud (ATLAS), “Measurement of dijet azimuthal decorrelations in pp collisions at $\sqrt{s} = 8 \text{ TeV}$ with the ATLAS detector and determination of the strong coupling”, Phys. Rev. D **98**, 092004 (2018).
- [302] M. Aaboud (ATLAS), “Measurement of inclusive jet and dijet cross-sections in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, JHEP **05**, 195 (2018).
- [303] M. Aaboud (ATLAS), “Measurement of the production cross-section of a single top quark in association with a Z boson in proton-proton collisions at 13 TeV with the ATLAS detector”, Phys. Lett. B **780**, 557–577 (2018).
- [304] M. Aaboud (ATLAS), “Observation of Higgs boson production in association with a top quark pair at the LHC with the ATLAS detector”, Phys. Lett. B **784**, 173–191 (2018).
- [305] M. Aaboud (ATLAS), “Search for a Structure in the $B_s^0\pi^\pm$ Invariant Mass Spectrum with the ATLAS Experiment”, Phys. Rev. Lett. **120**, 202007 (2018).
- [306] M. Aaboud (ATLAS), “Search for an invisibly decaying Higgs boson or dark matter candidates produced in association with a Z boson in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Phys. Lett. B **776**, 318–337 (2018).
- [307] M. Aaboud (ATLAS), “Search for dark matter in events with a hadronically decaying vector boson and missing transverse momentum in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, JHEP **10**, 180 (2018).
- [308] M. Aaboud (ATLAS), “Search for electroweak production of supersymmetric particles in final states with two or three leptons at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Eur. Phys. J. C **78**, 995 (2018).
- [309] M. Aaboud (ATLAS), “Search for exclusive Higgs and Z boson decays to $\phi\gamma$ and $\rho\gamma$ with the ATLAS detector”, JHEP **07**, 127 (2018).
- [310] M. Aaboud (ATLAS), “Search for flavour-changing neutral current top-quark decays $t \rightarrow qZ$ in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, JHEP **07**, 176 (2018).
- [311] M. Aaboud (ATLAS), “Search for heavy ZZ resonances in the $\ell^+\ell^-\ell^+\ell^-$ and $\ell^+\ell^-\nu\bar{\nu}$ final states using proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Eur. Phys. J. C **78**, 293 (2018).
- [312] M. Aaboud (ATLAS), “Search for Higgs boson pair production in the $\gamma\gamma b\bar{b}$ final state with 13 TeV pp collision data collected by the ATLAS experiment”, JHEP **11**, 040 (2018).
- [313] M. Aaboud (ATLAS), “Search for low-mass dijet resonances using trigger-level jets with the ATLAS detector in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ ”, Phys. Rev. Lett. **121**, 081801 (2018).
- [314] M. Aaboud (ATLAS), “Search for pair production of heavy vector-like quarks decaying into hadronic final states in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Phys. Rev. D **98**, 092005 (2018).
- [315] M. Aaboud (ATLAS), “Search for pair production of higgsinos in final states with at least three b -tagged jets in $\sqrt{s} = 13 \text{ TeV}$ pp collisions using the ATLAS detector”, Phys. Rev. D **98**, 092002 (2018).
- [316] M. Aaboud (ATLAS), “Search for resonant WZ production in the fully leptonic final state in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector”, Phys. Lett. B **787**, 68–88 (2018).
- [317] M. Aaboud (ATLAS), “Search for the Decay of the Higgs Boson to Charm Quarks with the ATLAS Experiment”, Phys. Rev. Lett. **120**, 211802 (2018).

- [318] M. Aaboud (ATLAS), “Search for the Higgs boson produced in association with a vector boson and decaying into two spin-zero particles in the $H \rightarrow aa \rightarrow 4b$ channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **10**, 031 (2018).
- [319] M. Aaboud (ATLAS), “ $ZZ \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$ cross-section measurements and search for anomalous triple gauge couplings in 13 TeV pp collisions with the ATLAS detector”, Phys. Rev. D **97**, 032005 (2018).
- [320] M. Aaboud (ATLAS), “A search for lepton-flavor-violating decays of the Z boson into a τ -lepton and a light lepton with the ATLAS detector”, Phys. Rev. D **98**, 092010 (2018).
- [321] M. Aaboud (ATLAS), “A search for pair-produced resonances in four-jet final states at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 250 (2018).
- [322] M. Aaboud (ATLAS), “A search for resonances decaying into a Higgs boson and a new particle X in the $XH \rightarrow qqbb$ final state with the ATLAS detector”, Phys. Lett. B **779**, 24–45 (2018).
- [323] M. Aaboud (ATLAS), “Angular analysis of $B_d^0 \rightarrow K^* \mu^+ \mu^-$ decays in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **10**, 047 (2018).
- [324] M. Aaboud (ATLAS, CMS), “Combination of inclusive and differential $t\bar{t}$ charge asymmetry measurements using ATLAS and CMS data at $\sqrt{s} = 7$ and 8 TeV”, JHEP **04**, 033 (2018).
- [325] M. Aaboud (ATLAS), “Combination of searches for heavy resonances decaying into bosonic and leptonic final states using 36 fb $^{-1}$ of proton-proton collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **98**, 052008 (2018).
- [326] M. Aaboud (ATLAS), “Combination of the searches for pair-produced vector-like partners of the third-generation quarks at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. Lett. **121**, 211801 (2018).
- [327] M. Aaboud (ATLAS), “Comparison between simulated and observed LHC beam backgrounds in the ATLAS experiment at $E_{\text{beam}} = 4$ TeV”, JINST **13**, P12006 (2018).
- [328] M. Aaboud (ATLAS), “Constraints on off-shell Higgs boson production and the Higgs boson total width in $ZZ \rightarrow 4\ell$ and $ZZ \rightarrow 2\ell 2\nu$ final states with the ATLAS detector”, Phys. Lett. B **786**, 223–244 (2018).
- [329] M. Aaboud (ATLAS), “Direct top-quark decay width measurement in the $t\bar{t}$ lepton+jets channel at $\sqrt{s}=8$ TeV with the ATLAS experiment”, Eur. Phys. J. C **78**, 129 (2018).
- [330] M. Aaboud (ATLAS), “Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector”, Phys. Rev. D **97**, 072003 (2018).
- [331] M. Aaboud (ATLAS), “Measurement of τ polarisation in $Z/\gamma^* \rightarrow \tau\tau$ decays in proton–proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 163 (2018).
- [332] M. Aaboud (ATLAS), “Measurement of colour flow using jet-pull observables in $t\bar{t}$ events with the ATLAS experiment at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **78**, 847 (2018).
- [333] M. Aaboud (ATLAS), “Measurement of differential cross sections and W^+/W^- cross-section ratios for W boson production in association with jets at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **05**, [Erratum: JHEP 10, 048 (2020)], 077 (2018).
- [334] M. Aaboud (ATLAS), “Measurement of differential cross sections of isolated-photon plus heavy-flavour jet production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Lett. B **776**, 295–317 (2018).
- [335] M. Aaboud (ATLAS), “Measurement of differential cross-sections of a single top quark produced in association with a W boson at $\sqrt{s} = 13$ TeV with ATLAS”, Eur. Phys. J. C **78**, 186 (2018).
- [336] M. Aaboud (ATLAS), “Measurement of jet fragmentation in 5.02 TeV proton-lead and proton-proton collisions with the ATLAS detector”, Nucl. Phys. A **978**, 65 (2018).
- [337] M. Aaboud (ATLAS), “Measurement of jet fragmentation in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, Phys. Rev. C **98**, 024908 (2018).
- [338] M. Aaboud (ATLAS), “Measurement of long-range multiparticle azimuthal correlations with the subevent cumulant method in pp and $p + Pb$ collisions with the ATLAS detector at the CERN Large Hadron Collider”, Phys. Rev. C **97**, 024904 (2018).
- [339] M. Aaboud (ATLAS), “Measurement of longitudinal flow decorrelations in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ and 5.02 TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 142 (2018).
- [340] M. Aaboud (ATLAS), “Measurement of quarkonium production in proton–lead and proton–proton collisions at 5.02 TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 171 (2018).
- [341] M. Aaboud (ATLAS), “Measurement of the $Z\gamma \rightarrow \nu\bar{\nu}\gamma$ production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector and limits on anomalous triple gauge-boson couplings”, JHEP **12**, 010 (2018).

- [342] M. Aaboud (ATLAS), “Measurement of the W -boson mass in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **78**, [Erratum: *Eur.Phys.J.C* 78, 898 (2018)], 110 (2018).
- [343] M. Aaboud (ATLAS), “Measurement of the azimuthal anisotropy of charged particles produced in $\sqrt{s_{NN}} = 5.02$ TeV Pb+Pb collisions with the ATLAS detector”, *Eur. Phys. J. C* **78**, 997 (2018).
- [344] M. Aaboud (ATLAS), “Measurement of the cross section for isolated-photon plus jet production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Lett. B* **780**, 578–602 (2018).
- [345] M. Aaboud (ATLAS), “Measurement of the cross-section for producing a W boson in association with a single top quark in pp collisions at $\sqrt{s} = 13$ TeV with ATLAS”, *JHEP* **01**, 063 (2018).
- [346] M. Aaboud (ATLAS), “Measurement of the exclusive $\gamma\gamma \rightarrow \mu^+\mu^-$ process in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **777**, 303–323 (2018).
- [347] M. Aaboud (ATLAS), “Measurement of the Higgs boson coupling properties in the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channel at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **03**, 095 (2018).
- [348] M. Aaboud (ATLAS), “Measurement of the Higgs boson mass in the $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ channels with $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector”, *Phys. Lett. B* **784**, 345–366 (2018).
- [349] M. Aaboud (ATLAS), “Measurement of the inclusive and fiducial $t\bar{t}$ production cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **78**, 487 (2018).
- [350] M. Aaboud (ATLAS), “Measurement of the production cross section of three isolated photons in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Phys. Lett. B* **781**, 55–76 (2018).
- [351] M. Aaboud (ATLAS), “Measurement of the Soft-Drop Jet Mass in pp Collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **121**, 092001 (2018).
- [352] M. Aaboud (ATLAS), “Measurement of the suppression and azimuthal anisotropy of muons from heavy-flavor decays in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, *Phys. Rev. C* **98**, 044905 (2018).
- [353] M. Aaboud (ATLAS), “Measurements of $t\bar{t}$ differential cross-sections of highly boosted top quarks decaying to all-hadronic final states in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Rev. D* **98**, 012003 (2018).
- [354] M. Aaboud (ATLAS), “Measurements of b-jet tagging efficiency with the ATLAS detector using $t\bar{t}$ events at $\sqrt{s} = 13$ TeV”, *JHEP* **08**, 089 (2018).
- [355] M. Aaboud (ATLAS), “Measurements of differential cross sections of top quark pair production in association with jets in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *JHEP* **10**, 159 (2018).
- [356] M. Aaboud (ATLAS), “Measurements of Higgs boson properties in the diphoton decay channel with 36 fb⁻¹ of pp collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **98**, 052005 (2018).
- [357] M. Aaboud (ATLAS), “Observation of $H \rightarrow b\bar{b}$ decays and VH production with the ATLAS detector”, *Phys. Lett. B* **786**, 59–86 (2018).
- [358] M. Aaboud (ATLAS), “Observation of centrality-dependent acoplanarity for muon pairs produced via two-photon scattering in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **121**, 212301 (2018).
- [359] M. Aaboud (ATLAS), “Operation and performance of the ATLAS Tile Calorimeter in Run 1”, *Eur. Phys. J. C* **78**, 987 (2018).
- [360] M. Aaboud (ATLAS), “Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **78**, 903 (2018).
- [361] M. Aaboud (ATLAS), “Probing the quantum interference between singly and doubly resonant top-quark production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **121**, 152002 (2018).
- [362] M. Aaboud (ATLAS), “Prompt and non-prompt J/ψ and $\psi(2S)$ suppression at high transverse momentum in 5.02 TeV Pb+Pb collisions with the ATLAS experiment”, *Eur. Phys. J. C* **78**, 762 (2018).
- [363] M. Aaboud (ATLAS), “Prompt and non-prompt J/ψ elliptic flow in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **78**, 784 (2018).
- [364] M. Aaboud (ATLAS), “Search for $W' \rightarrow t\bar{b}$ decays in the hadronic final state using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **781**, 327–348 (2018).
- [365] M. Aaboud (ATLAS), “Search for WW/WZ resonance production in $\ell\nu qq$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **03**, 042 (2018).

- [366] M. Aaboud (ATLAS), “Search for a heavy Higgs boson decaying into a Z boson and another heavy Higgs boson in the $\ell\ell b\bar{b}$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **783**, 392–414 (2018).
- [367] M. Aaboud (ATLAS), “Search for a new heavy gauge boson resonance decaying into a lepton and missing transverse momentum in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, Eur. Phys. J. C **78**, 401 (2018).
- [368] M. Aaboud (ATLAS), “Search for additional heavy neutral Higgs and gauge bosons in the ditau final state produced in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **01**, 055 (2018).
- [369] M. Aaboud (ATLAS), “Search for B-L R -parity-violating top squarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS experiment”, Phys. Rev. D **97**, 032003 (2018).
- [370] M. Aaboud (ATLAS), “Search for charged Higgs bosons decaying into top and bottom quarks at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **11**, 085 (2018).
- [371] M. Aaboud (ATLAS), “Search for charged Higgs bosons decaying via $H^\pm \rightarrow \tau^\pm \nu_\tau$ in the τ +jets and τ +lepton final states with 36 fb^{-1} of pp collision data recorded at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, JHEP **09**, 139 (2018).
- [372] M. Aaboud (ATLAS), “Search for chargino-neutralino production using recursive jigsaw reconstruction in final states with two or three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **98**, 092012 (2018).
- [373] M. Aaboud (ATLAS), “Search for dark matter and other new phenomena in events with an energetic jet and large missing transverse momentum using the ATLAS detector”, JHEP **01**, 126 (2018).
- [374] M. Aaboud (ATLAS), “Search for dark matter produced in association with bottom or top quarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **78**, 18 (2018).
- [375] M. Aaboud (ATLAS), “Search for diboson resonances with boson-tagged jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **777**, 91–113 (2018).
- [376] M. Aaboud (ATLAS), “Search for doubly charged Higgs boson production in multi-lepton final states with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **78**, 199 (2018).
- [377] M. Aaboud (ATLAS), “Search for electroweak production of supersymmetric states in scenarios with compressed mass spectra at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **97**, 052010 (2018).
- [378] M. Aaboud (ATLAS), “Search for flavor-changing neutral currents in top quark decays $t \rightarrow Hc$ and $t \rightarrow Hu$ in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **98**, 032002 (2018).
- [379] M. Aaboud (ATLAS), “Search for heavy particles decaying into top-quark pairs using lepton-plus-jets events in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 565 (2018).
- [380] M. Aaboud (ATLAS), “Search for heavy resonances decaying into WW in the $e\nu\mu\nu$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 24 (2018).
- [381] M. Aaboud (ATLAS), “Search for heavy resonances decaying into a W or Z boson and a Higgs boson in final states with leptons and b -jets in 36 fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, JHEP **03**, [Erratum: JHEP 11, 051 (2018)], 174 (2018).
- [382] M. Aaboud (ATLAS), “Search for heavy resonances decaying to a photon and a hadronically decaying $Z/W/H$ boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **98**, 032015 (2018).
- [383] M. Aaboud (ATLAS), “Search for Higgs boson decays into pairs of light (pseudo)scalar particles in the $\gamma\gamma jj$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **782**, 750–767 (2018).
- [384] M. Aaboud (ATLAS), “Search for Higgs boson decays to beyond-the-Standard-Model light bosons in four-lepton events with the ATLAS detector at $\sqrt{s} = 13$ TeV”, JHEP **06**, 166 (2018).
- [385] M. Aaboud (ATLAS), “Search for Higgs boson pair production in the $\gamma\gamma WW^*$ channel using pp collision data recorded at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **78**, 1007 (2018).
- [386] M. Aaboud (ATLAS), “Search for Higgs bosons produced via vector-boson fusion and decaying into bottom quark pairs in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Phys. Rev. D **98**, 052003 (2018).
- [387] M. Aaboud (ATLAS), “Search for High-Mass Resonances Decaying to $\tau\nu$ in pp Collisions at $\sqrt{s}=13$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **120**, 161802 (2018).

- [388] M. Aaboud (ATLAS), “Search for lepton-flavor violation in different-flavor, high-mass final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **98**, 092008 (2018).
- [389] M. Aaboud (ATLAS), “Search for long-lived charginos based on a disappearing-track signature in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **06**, 022 (2018).
- [390] M. Aaboud (ATLAS), “Search for long-lived, massive particles in events with displaced vertices and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **97**, 052012 (2018).
- [391] M. Aaboud (ATLAS), “Search for new phenomena in events with same-charge leptons and b -jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **12**, 039 (2018).
- [392] M. Aaboud (ATLAS), “Search for new phenomena in high-mass final states with a photon and a jet from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **78**, 102 (2018).
- [393] M. Aaboud (ATLAS), “Search for new phenomena using the invariant mass distribution of same-flavour opposite-sign dilepton pairs in events with missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **78**, 625 (2018).
- [394] M. Aaboud (ATLAS), “Search for pair production of heavy vector-like quarks decaying into high- p_T W bosons and top quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **08**, 048 (2018).
- [395] M. Aaboud (ATLAS), “Search for pair production of up-type vector-like quarks and for four-top-quark events in final states with multiple b -jets with the ATLAS detector”, *JHEP* **07**, 089 (2018).
- [396] M. Aaboud (ATLAS), “Search for pair- and single-production of vector-like quarks in final states with at least one Z boson decaying into a pair of electrons or muons in pp collision data collected with the ATLAS detector at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **98**, 112010 (2018).
- [397] M. Aaboud (ATLAS), “Search for photonic signatures of gauge-mediated supersymmetry in 13 TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **97**, 092006 (2018).
- [398] M. Aaboud (ATLAS), “Search for R-parity-violating supersymmetric particles in multi-jet final states produced in p - p collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC”, *Phys. Lett. B* **785**, 136–158 (2018).
- [399] M. Aaboud (ATLAS), “Search for resonances in the mass distribution of jet pairs with one or two jets identified as b -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **98**, 032016 (2018).
- [400] M. Aaboud (ATLAS), “Search for resonant and non-resonant Higgs boson pair production in the $b\bar{b}\tau^+\tau^-$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **121**, [Erratum: *Phys.Rev.Lett.* 122, 089901 (2019)], 191801 (2018).
- [401] M. Aaboud (ATLAS), “Search for squarks and gluinos in final states with jets and missing transverse momentum using 36 fb $^{-1}$ of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, *Phys. Rev. D* **97**, 112001 (2018).
- [402] M. Aaboud (ATLAS), “Search for supersymmetry in events with four or more leptons in $\sqrt{s} = 13$ TeV pp collisions with ATLAS”, *Phys. Rev. D* **98**, 032009 (2018).
- [403] M. Aaboud (ATLAS), “Search for supersymmetry in final states with charm jets and missing transverse momentum in 13 TeV pp collisions with the ATLAS detector”, *JHEP* **09**, 050 (2018).
- [404] M. Aaboud (ATLAS), “Search for supersymmetry in final states with missing transverse momentum and multiple b -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **06**, 107 (2018).
- [405] M. Aaboud (ATLAS), “Search for the direct production of charginos and neutralinos in final states with tau leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **78**, 154 (2018).
- [406] M. Aaboud (ATLAS), “Search for the standard model Higgs boson produced in association with top quarks and decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **97**, 072016 (2018).
- [407] M. Aaboud (ATLAS), “Search for top squarks decaying to tau sleptons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **98**, 032008 (2018).
- [408] M. Aaboud (ATLAS), “Search for top-squark pair production in final states with one lepton, jets, and missing transverse momentum using 36 fb $^{-1}$ of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, *JHEP* **06**, 108 (2018).
- [409] M. Aaboud (ATLAS), “Searches for exclusive Higgs and Z boson decays into $J/\psi\gamma$, $\psi(2S)\gamma$, and $\Upsilon(nS)\gamma$ at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **786**, 134–155 (2018).

- [410] M. Aaboud (ATLAS), “Searches for heavy ZZ and ZW resonances in the $\ell\ell q\bar{q}$ and $\nu\nu q\bar{q}$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **03**, 009 (2018).
- [411] K. Albertsson, “Machine Learning in High Energy Physics Community White Paper”, *J. Phys. Conf. Ser.* **1085**, 022008 (2018).
- [412] L. Bauerdick (HEP Software Foundation), “HEP Software Foundation Community White Paper Working Group - Data Analysis and Interpretation”, edited by M. S. Neubauer (2018).
- [413] M. Bellis, “HEP Software Foundation Community White Paper Working Group – Visualization”, (2018).
- [414] D. Berzano (HEP Software Foundation), “HEP Software Foundation Community White Paper Working Group - Training, Staffing and Careers”, (2018).
- [415] M. D. Hildreth, “HEP Software Foundation Community White Paper Working Group - Data and Software Preservation to Enable Reuse”, (2018).
- [416] M. Aaboud (ATLAS), “Evidence for the $H \rightarrow b\bar{b}$ decay with the ATLAS detector”, *JHEP* **12**, 024 (2017).
- [417] M. Aaboud (ATLAS), “Jet energy scale measurements and their systematic uncertainties in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **96**, 072002 (2017).
- [418] M. Aaboud (ATLAS), “Measurement of b -hadron pair production with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **11**, 062 (2017).
- [419] M. Aaboud (ATLAS), “Measurement of lepton differential distributions and the top quark mass in $t\bar{t}$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77**, 804 (2017).
- [420] M. Aaboud (ATLAS), “Measurement of the cross-section for electroweak production of dijets in association with a Z boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **775**, 206–228 (2017).
- [421] M. Aaboud (ATLAS), “Measurement of the Drell-Yan triple-differential cross section in pp collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **12**, 059 (2017).
- [422] M. Aaboud (ATLAS), “Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *JHEP* **11**, 191 (2017).
- [423] M. Aaboud (ATLAS), “Performance of the ATLAS Track Reconstruction Algorithms in Dense Environments in LHC Run 2”, *Eur. Phys. J. C* **77**, 673 (2017).
- [424] M. Aaboud (ATLAS), “Search for direct top squark pair production in final states with two leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **77**, 898 (2017).
- [425] M. Aaboud (ATLAS), “Search for pair production of heavy vector-like quarks decaying to high- p_T W bosons and b quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **10**, 141 (2017).
- [426] M. Aaboud (ATLAS), “Searches for the $Z\gamma$ decay mode of the Higgs boson and for new high-mass resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **10**, 112 (2017).
- [427] M. Aaboud (ATLAS), “Study of $WW\gamma$ and $WZ\gamma$ production in pp collisions at $\sqrt{s} = 8$ TeV and search for anomalous quartic gauge couplings with the ATLAS experiment”, *Eur. Phys. J. C* **77**, 646 (2017).
- [428] M. Aaboud (ATLAS), “Study of ordered hadron chains with the ATLAS detector”, *Phys. Rev. D* **96**, 092008 (2017).
- [429] M. Aaboud (ATLAS), “A measurement of the calorimeter response to single hadrons and determination of the jet energy scale uncertainty using LHC Run-1 pp -collision data with the ATLAS detector”, *Eur. Phys. J. C* **77**, 26 (2017).
- [430] M. Aaboud (ATLAS), “Analysis of the Wtb vertex from the measurement of triple-differential angular decay rates of single top quarks produced in the t -channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **12**, 017 (2017).
- [431] M. Aaboud (ATLAS), “Determination of the strong coupling constant α_s from transverse energy–energy correlations in multijet events at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **77**, 872 (2017).
- [432] M. Aaboud (ATLAS), “Electron efficiency measurements with the ATLAS detector using 2012 LHC proton–proton collision data”, *Eur. Phys. J. C* **77**, 195 (2017).
- [433] M. Aaboud (ATLAS), “Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC”, *Nature Phys.* **13**, 852–858 (2017).
- [434] M. Aaboud (ATLAS), “Femtoscopia with identified charged pions in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ATLAS”, *Phys. Rev. C* **96**, 064908 (2017).

- [435] M. Aaboud (ATLAS), “Fiducial, total and differential cross-section measurements of t -channel single top-quark production in pp collisions at 8 TeV using data collected by the ATLAS detector”, *Eur. Phys. J. C* **77**, 531 (2017).
- [436] M. Aaboud (ATLAS), “High- E_T isolated-photon plus jets production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Nucl. Phys. B* **918**, 257–316 (2017).
- [437] M. Aaboud (ATLAS), “Identification and rejection of pile-up jets at high pseudorapidity with the ATLAS detector”, *Eur. Phys. J. C* **77**, [Erratum: *Eur. Phys. J. C* **77**, 712 (2017)], 580 (2017).
- [438] M. Aaboud (ATLAS), “Jet reconstruction and performance using particle flow with the ATLAS Detector”, *Eur. Phys. J. C* **77**, 466 (2017).
- [439] M. Aaboud (ATLAS), “Measurement of $W^\pm W^\pm$ vector-boson scattering and limits on anomalous quartic gauge couplings with the ATLAS detector”, *Phys. Rev. D* **96**, 012007 (2017).
- [440] M. Aaboud (ATLAS), “Measurement of W boson angular distributions in events with high transverse momentum jets at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Phys. Lett. B* **765**, 132–153 (2017).
- [441] M. Aaboud (ATLAS), “Measurement of $WW/WZ \rightarrow \ell\nu qq'$ production with the hadronically decaying boson reconstructed as one or two jets in pp collisions at $\sqrt{s} = 8$ TeV with ATLAS, and constraints on anomalous gauge couplings”, *Eur. Phys. J. C* **77**, 563 (2017).
- [442] M. Aaboud (ATLAS), “Measurement of charged-particle distributions sensitive to the underlying event in $\sqrt{s} = 13$ TeV proton-proton collisions with the ATLAS detector at the LHC”, *JHEP* **03**, 157 (2017).
- [443] M. Aaboud (ATLAS), “Measurement of detector-corrected observables sensitive to the anomalous production of events with jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **77**, 765 (2017).
- [444] M. Aaboud (ATLAS), “Measurement of forward-backward multiplicity correlations in lead-lead, proton-lead, and proton-proton collisions with the ATLAS detector”, *Phys. Rev. C* **95**, 064914 (2017).
- [445] M. Aaboud (ATLAS), “Measurement of inclusive and differential cross sections in the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **10**, 132 (2017).
- [446] M. Aaboud (ATLAS), “Measurement of jet p_T correlations in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, *Phys. Lett. B* **774**, 379–402 (2017).
- [447] M. Aaboud (ATLAS), “Measurement of jet activity produced in top-quark events with an electron, a muon and two b -tagged jets in the final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77**, 220 (2017).
- [448] M. Aaboud (ATLAS), “Measurement of jet fragmentation in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC”, *Eur. Phys. J. C* **77**, 379 (2017).
- [449] M. Aaboud (ATLAS), “Measurement of multi-particle azimuthal correlations in pp , p +Pb and low-multiplicity Pb+Pb collisions with the ATLAS detector”, *Eur. Phys. J. C* **77**, 428 (2017).
- [450] M. Aaboud (ATLAS), “Measurement of the $t\bar{t}\gamma$ production cross section in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **11**, 086 (2017).
- [451] M. Aaboud (ATLAS), “Measurement of the k_t splitting scales in $Z \rightarrow \ell\ell$ events in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **08**, 026 (2017).
- [452] M. Aaboud (ATLAS), “Measurement of the $t\bar{t}$ production cross section in the τ + jets final state in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Phys. Rev. D* **95**, 072003 (2017).
- [453] M. Aaboud (ATLAS), “Measurement of the $t\bar{t}Z$ and $t\bar{t}W$ production cross sections in multilepton final states using 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77**, 40 (2017).
- [454] M. Aaboud (ATLAS), “Measurement of the W^+W^- production cross section in pp collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV with the ATLAS experiment”, *Phys. Lett. B* **773**, 354–374 (2017).
- [455] M. Aaboud (ATLAS), “Measurement of the ZZ production cross section in proton-proton collisions at $\sqrt{s} = 8$ TeV using the $ZZ \rightarrow \ell^-\ell^+\ell'^-\ell'^+$ and $ZZ \rightarrow \ell^-\ell^+\nu\bar{\nu}$ channels with the ATLAS detector”, *JHEP* **01**, 099 (2017).
- [456] M. Aaboud (ATLAS), “Measurement of the cross section for inclusive isolated-photon production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Lett. B* **770**, 473–493 (2017).
- [457] M. Aaboud (ATLAS), “Measurement of the inclusive cross-sections of single top-quark and top-antiquark t -channel production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **04**, 086 (2017).
- [458] M. Aaboud (ATLAS), “Measurement of the inclusive jet cross-sections in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **09**, 020 (2017).

- [459] M. Aaboud (ATLAS), “Measurement of the prompt J/ψ pair production cross-section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77**, 76 (2017).
- [460] M. Aaboud (ATLAS), “Measurement of the W boson polarisation in $t\bar{t}$ events from pp collisions at $\sqrt{s} = 8$ TeV in the lepton + jets channel with ATLAS”, *Eur. Phys. J. C* **77**, [Erratum: *Eur.Phys.J.C* 79, 19 (2019)], 264 (2017).
- [461] M. Aaboud (ATLAS), “Measurements of $\psi(2S)$ and $X(3872) \rightarrow J/\psi\pi^+\pi^-$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **01**, 117 (2017).
- [462] M. Aaboud (ATLAS), “Measurements of charge and CP asymmetries in b -hadron decays using top-quark events collected by the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **02**, 071 (2017).
- [463] M. Aaboud (ATLAS), “Measurements of electroweak Wjj production and constraints on anomalous gauge couplings with the ATLAS detector”, *Eur. Phys. J. C* **77**, 474 (2017).
- [464] M. Aaboud (ATLAS), “Measurements of integrated and differential cross sections for isolated photon pair production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **95**, 112005 (2017).
- [465] M. Aaboud (ATLAS), “Measurements of long-range azimuthal anisotropies and associated Fourier coefficients for pp collisions at $\sqrt{s} = 5.02$ and 13 TeV and p +Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Phys. Rev. C* **96**, 024908 (2017).
- [466] M. Aaboud (ATLAS), “Measurements of the production cross section of a Z boson in association with jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77**, 361 (2017).
- [467] M. Aaboud (ATLAS), “Measurements of top quark spin observables in $t\bar{t}$ events using dilepton final states in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *JHEP* **03**, 113 (2017).
- [468] M. Aaboud (ATLAS), “Measurements of top-quark pair differential cross-sections in the $e\mu$ channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **77**, 292 (2017).
- [469] M. Aaboud (ATLAS), “Measurements of top-quark pair to Z -boson cross-section ratios at $\sqrt{s} = 13, 8, 7$ TeV with the ATLAS detector”, *JHEP* **02**, 117 (2017).
- [470] M. Aaboud (ATLAS), “Performance of the ATLAS Transition Radiation Tracker in Run 1 of the LHC: tracker properties”, *JINST* **12**, P05002 (2017).
- [471] M. Aaboud (ATLAS), “Performance of the ATLAS Trigger System in 2015”, *Eur. Phys. J. C* **77**, 317 (2017).
- [472] M. Aaboud (ATLAS), “Precision measurement and interpretation of inclusive W^+ , W^- and Z/γ^* production cross sections with the ATLAS detector”, *Eur. Phys. J. C* **77**, 367 (2017).
- [473] M. Aaboud (ATLAS), “Probing the W tb vertex structure in t-channel single-top-quark production and decay in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **04**, 124 (2017).
- [474] M. Aaboud (ATLAS), “Reconstruction of primary vertices at the ATLAS experiment in Run 1 proton–proton collisions at the LHC”, *Eur. Phys. J. C* **77**, 332 (2017).
- [475] M. Aaboud (ATLAS), “Search for a scalar partner of the top quark in the jets plus missing transverse momentum final state at $\sqrt{s}=13$ TeV with the ATLAS detector”, *JHEP* **12**, 085 (2017).
- [476] M. Aaboud (ATLAS), “Search for anomalous electroweak production of WW/WZ in association with a high-mass dijet system in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **95**, 032001 (2017).
- [477] M. Aaboud (ATLAS), “Search for dark matter at $\sqrt{s} = 13$ TeV in final states containing an energetic photon and large missing transverse momentum with the ATLAS detector”, *Eur. Phys. J. C* **77**, 393 (2017).
- [478] M. Aaboud (ATLAS), “Search for dark matter in association with a Higgs boson decaying to b -quarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **765**, 11–31 (2017).
- [479] M. Aaboud (ATLAS), “Search for dark matter in association with a Higgs boson decaying to two photons at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **96**, 112004 (2017).
- [480] M. Aaboud (ATLAS), “Search for Dark Matter Produced in Association with a Higgs Boson Decaying to $b\bar{b}$ using 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **119**, 181804 (2017).
- [481] M. Aaboud (ATLAS), “Search for direct top squark pair production in events with a Higgs or Z boson, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *JHEP* **08**, 006 (2017).
- [482] M. Aaboud (ATLAS), “Search for Heavy Higgs Bosons A/H Decaying to a Top Quark Pair in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **119**, 191803 (2017).

- [483] M. Aaboud (ATLAS), “Search for heavy resonances decaying to a W or Z boson and a Higgs boson in the $q\bar{q}^{(\prime)}b\bar{b}$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **774**, 494–515 (2017).
- [484] M. Aaboud (ATLAS), “Search for heavy resonances decaying to a Z boson and a photon in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **764**, 11–30 (2017).
- [485] M. Aaboud (ATLAS), “Search for new high-mass phenomena in the dilepton final state using 36 fb¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **10**, 182 (2017).
- [486] M. Aaboud (ATLAS), “Search for new phenomena in a lepton plus high jet multiplicity final state with the ATLAS experiment using $\sqrt{s} = 13$ TeV proton-proton collision data”, JHEP **09**, 088 (2017).
- [487] M. Aaboud (ATLAS), “Search for new phenomena in dijet events using 37 fb⁻¹ of pp collision data collected at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **96**, 052004 (2017).
- [488] M. Aaboud (ATLAS), “Search for new phenomena in events containing a same-flavour opposite-sign dilepton pair, jets, and large missing transverse momentum in $\sqrt{s} = 13$ pp collisions with the ATLAS detector”, Eur. Phys. J. C **77**, 144 (2017).
- [489] M. Aaboud (ATLAS), “Search for new phenomena in high-mass diphoton final states using 37 fb⁻¹ of proton-proton collisions collected at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **775**, 105–125 (2017).
- [490] M. Aaboud (ATLAS), “Search for new phenomena with large jet multiplicities and missing transverse momentum using large-radius jets and flavour-tagging at ATLAS in 13 TeV pp collisions”, JHEP **12**, 034 (2017).
- [491] M. Aaboud (ATLAS), “Search for new resonances decaying to a W or Z boson and a Higgs boson in the $\ell^+\ell^-\bar{b}b$, $\ell\nu b\bar{b}$, and $\nu\bar{\nu}b\bar{b}$ channels with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **765**, 32–52 (2017).
- [492] M. Aaboud (ATLAS), “Search for pair production of vector-like top quarks in events with one lepton, jets, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, JHEP **08**, 052 (2017).
- [493] M. Aaboud (ATLAS), “Search for squarks and gluinos in events with an isolated lepton, jets, and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **96**, 112010 (2017).
- [494] M. Aaboud (ATLAS), “Search for supersymmetry in events with b -tagged jets and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **11**, 195 (2017).
- [495] M. Aaboud (ATLAS), “Search for supersymmetry in final states with two same-sign or three leptons and jets using 36 fb⁻¹ of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, JHEP **09**, [Erratum: JHEP 08, 121 (2019)], 084 (2017).
- [496] M. Aaboud (ATLAS), “Search for the dimuon decay of the Higgs boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. Lett. **119**, 051802 (2017).
- [497] M. Aaboud (ATLAS), “Search for top quark decays $t \rightarrow qH$, with $H \rightarrow \gamma\gamma$, in $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector”, JHEP **10**, 129 (2017).
- [498] M. Aaboud (ATLAS), “Search for triboson $W^\pm W^\pm W^\mp$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **77**, 141 (2017).
- [499] M. Aaboud (ATLAS), “Studies of $Z\gamma$ production in association with a high-mass dijet system in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **07**, 107 (2017).
- [500] M. Aaboud (ATLAS), “Study of the material of the ATLAS inner detector for Run 2 of the LHC”, JINST **12**, P12009 (2017).
- [501] M. Aaboud (ATLAS), “Top-quark mass measurement in the all-hadronic $t\bar{t}$ decay channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 118 (2017).
- [502] G. Aad (ATLAS), “Performance of algorithms that reconstruct missing transverse momentum in $\sqrt{s} = 8$ TeV proton-proton collisions in the ATLAS detector”, Eur. Phys. J. C **77**, 241 (2017).
- [503] G. Aad (ATLAS), “Search for lepton-flavour-violating decays of the Higgs and Z bosons with the ATLAS detector”, Eur. Phys. J. C **77**, 70 (2017).
- [504] G. Aad (ATLAS), “Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1”, Eur. Phys. J. C **77**, 490 (2017).
- [505] T. Aaltonen (CDF), “Observation of the $Y(4140)$ Structure in the $J/\psi\phi$ Mass Spectrum in $B^\pm \rightarrow J/\psi\phi K^\pm$ Decays”, Mod. Phys. Lett. A **32**, 1750139 (2017).
- [506] P. Elmer, **M. Neubauer**, M. D. Sokoloff, “Strategic Plan for a Scientific Software Innovation Institute (S2I2) for High Energy Physics”, (2017).

- [507] M. Aaboud (ATLAS), “Measurement of the Inelastic Proton-Proton Cross Section at $\sqrt{s} = 13$ TeV with the ATLAS Detector at the LHC”, Phys. Rev. Lett. **117**, 182002 (2016).
- [508] M. Aaboud (ATLAS), “A measurement of material in the ATLAS tracker using secondary hadronic interactions in 7 TeV pp collisions”, JINST **11**, P11020 (2016).
- [509] M. Aaboud (ATLAS), “Charged-particle distributions at low transverse momentum in $\sqrt{s} = 13$ TeV pp interactions measured with the ATLAS detector at the LHC”, Eur. Phys. J. C **76**, 502 (2016).
- [510] M. Aaboud (ATLAS), “Dark matter interpretations of ATLAS searches for the electroweak production of supersymmetric particles in $\sqrt{s} = 8$ TeV proton-proton collisions”, JHEP **09**, 175 (2016).
- [511] M. Aaboud (ATLAS), “Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC”, Eur. Phys. J. C **76**, 653 (2016).
- [512] M. Aaboud (ATLAS), “Measurement of W^+W^- production in association with one jet in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B **763**, 114–133 (2016).
- [513] M. Aaboud (ATLAS), “Measurement of exclusive $\gamma\gamma \rightarrow W^+W^-$ production and search for exclusive Higgs boson production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Rev. D **94**, 032011 (2016).
- [514] M. Aaboud (ATLAS), “Measurement of jet activity in top quark events using the $e\mu$ final state with two b -tagged jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 074 (2016).
- [515] M. Aaboud (ATLAS), “Measurement of the $b\bar{b}$ dijet cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 670 (2016).
- [516] M. Aaboud (ATLAS), “Measurement of the $t\bar{t}$ production cross-section using $e\mu$ events with b -tagged jets in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector”, Phys. Lett. B **761**, [Erratum: Phys.Lett.B 772, 879–879 (2017)], 136–157 (2016).
- [517] M. Aaboud (ATLAS), “Measurement of the $W^\pm Z$ boson pair-production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, Phys. Lett. B **762**, 1–22 (2016).
- [518] M. Aaboud (ATLAS), “Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run-1 data”, Eur. Phys. J. C **76**, 666 (2016).
- [519] M. Aaboud (ATLAS), “Measurement of the relative width difference of the $B^0-\bar{B}^0$ system with the ATLAS detector”, JHEP **06**, 081 (2016).
- [520] M. Aaboud (ATLAS), “Measurement of the top quark mass in the $t\bar{t} \rightarrow$ dilepton channel from $\sqrt{s} = 8$ TeV ATLAS data”, Phys. Lett. B **761**, 350–371 (2016).
- [521] M. Aaboud (ATLAS), “Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B **761**, 158–178 (2016).
- [522] M. Aaboud (ATLAS), “Measurement of top quark pair differential cross-sections in the dilepton channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS”, Phys. Rev. D **94**, [Addendum: Phys.Rev.D 101, 119901 (2020)], 092003 (2016).
- [523] M. Aaboud (ATLAS), “Search for bottom squark pair production in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 547 (2016).
- [524] M. Aaboud (ATLAS), “Search for charged Higgs bosons produced in association with a top quark and decaying via $H^\pm \rightarrow \tau\nu$ using pp collision data recorded at $\sqrt{s} = 13$ TeV by the ATLAS detector”, Phys. Lett. B **759**, 555–574 (2016).
- [525] M. Aaboud (ATLAS), “Search for dark matter produced in association with a hadronically decaying vector boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **763**, 251–268 (2016).
- [526] M. Aaboud (ATLAS), “Search for heavy long-lived charged R -hadrons with the ATLAS detector in 3.2 fb $^{-1}$ of proton-proton collision data at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **760**, 647–665 (2016).
- [527] M. Aaboud (ATLAS), “Search for Higgs and Z Boson Decays to $\phi\gamma$ with the ATLAS Detector”, Phys. Rev. Lett. **117**, 111802 (2016).
- [528] M. Aaboud (ATLAS), “Search for high-mass new phenomena in the dilepton final state using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **761**, 372–392 (2016).
- [529] M. Aaboud (ATLAS), “Search for metastable heavy charged particles with large ionization energy loss in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS experiment”, Phys. Rev. D **93**, 112015 (2016).
- [530] M. Aaboud (ATLAS), “Search for Minimal Supersymmetric Standard Model Higgs bosons H/A and for a Z' boson in the $\tau\tau$ final state produced in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, Eur. Phys. J. C **76**, 585 (2016).

- [531] M. Aaboud (ATLAS), “Search for new phenomena in different-flavour high-mass dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, 541 (2016).
- [532] M. Aaboud (ATLAS), “Search for new phenomena in events with a photon and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **06**, 059 (2016).
- [533] M. Aaboud (ATLAS), “Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Rev. D* **94**, 032005 (2016).
- [534] M. Aaboud (ATLAS), “Search for new resonances in events with one lepton and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **762**, 334–352 (2016).
- [535] M. Aaboud (ATLAS), “Search for pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **94**, 052002 (2016).
- [536] M. Aaboud (ATLAS), “Search for resonances in diphoton events at $\sqrt{s}=13$ TeV with the ATLAS detector”, *JHEP* **09**, 001 (2016).
- [537] M. Aaboud (ATLAS), “Search for resonances in the mass distribution of jet pairs with one or two jets identified as b -jets in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **759**, 229–246 (2016).
- [538] M. Aaboud (ATLAS), “Search for scalar leptoquarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, *New J. Phys.* **18**, 093016 (2016).
- [539] M. Aaboud (ATLAS), “Search for squarks and gluinos in events with hadronically decaying tau leptons, jets and missing transverse momentum in proton–proton collisions at $\sqrt{s} = 13$ TeV recorded with the ATLAS detector”, *Eur. Phys. J. C* **76**, 683 (2016).
- [540] M. Aaboud (ATLAS), “Search for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, 392 (2016).
- [541] M. Aaboud (ATLAS), “Search for supersymmetry in a final state containing two photons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions at the LHC using the ATLAS detector”, *Eur. Phys. J. C* **76**, 517 (2016).
- [542] M. Aaboud (ATLAS), “Search for TeV-scale gravity signatures in high-mass final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **760**, 520–537 (2016).
- [543] M. Aaboud (ATLAS), “Search for the Higgs boson produced in association with a W boson and decaying to four b -quarks via two spin-zero particles in pp collisions at 13 TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, 605 (2016).
- [544] M. Aaboud (ATLAS), “Search for the Standard Model Higgs boson produced by vector-boson fusion and decaying to bottom quarks in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *JHEP* **11**, 112 (2016).
- [545] M. Aaboud (ATLAS), “Search for top squarks in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **94**, 052009 (2016).
- [546] M. Aaboud (ATLAS), “Searches for heavy diboson resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **09**, 173 (2016).
- [547] M. Aaboud (ATLAS), “Study of hard double-parton scattering in four-jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, *JHEP* **11**, 110 (2016).
- [548] M. Aaboud (ATLAS), “Study of the rare decays of B_s^0 and B^0 into muon pairs from data collected during the LHC Run 1 with the ATLAS detector”, *Eur. Phys. J. C* **76**, 513 (2016).
- [549] G. Aad (ATLAS), “Search for single top-quark production via flavour-changing neutral currents at 8 TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, [Erratum: *Eur.Phys.J.C* 82, 70 (2022)], 55 (2016).
- [550] G. Aad (ATLAS), “A new method to distinguish hadronically decaying boosted Z bosons from W bosons using the ATLAS detector”, *Eur. Phys. J. C* **76**, 238 (2016).
- [551] G. Aad (ATLAS), “A search for an excited muon decaying to a muon and two jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *New J. Phys.* **18**, [Erratum: *New J.Phys.* 21, 109501 (2019)], 073021 (2016).
- [552] G. Aad (ATLAS), “A search for prompt lepton-jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **02**, 062 (2016).
- [553] G. Aad (ATLAS), “A search for top squarks with R-parity-violating decays to all-hadronic final states with the ATLAS detector in $\sqrt{s} = 8$ TeV proton-proton collisions”, *JHEP* **06**, 067 (2016).

- [554] G. Aad (ATLAS), “Beam-induced and cosmic-ray backgrounds observed in the ATLAS detector during the LHC 2012 proton-proton running period”, *JINST* **11**, P05013 (2016).
- [555] G. Aad (ATLAS), “Centrality, rapidity and transverse momentum dependence of isolated prompt photon production in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV measured with the ATLAS detector”, *Phys. Rev. C* **93**, 034914 (2016).
- [556] G. Aad (ATLAS), “Charged-particle distributions in pp interactions at $\sqrt{s} = 8$ TeV measured with the ATLAS detector”, *Eur. Phys. J. C* **76**, 403 (2016).
- [557] G. Aad (ATLAS), “Charged-particle distributions in $\sqrt{s} = 13$ TeV pp interactions measured with the ATLAS detector at the LHC”, *Phys. Lett. B* **758**, 67–88 (2016).
- [558] G. Aad (ATLAS), “Combination of searches for WW , WZ , and ZZ resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **755**, 285–305 (2016).
- [559] G. Aad (ATLAS), “Constraints on non-Standard Model Higgs boson interactions in an effective Lagrangian using differential cross sections measured in the $H \rightarrow \gamma\gamma$ decay channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **753**, 69–85 (2016).
- [560] G. Aad (ATLAS), “Dijet production in $\sqrt{s} = 7$ TeV pp collisions with large rapidity gaps at the ATLAS experiment”, *Phys. Lett. B* **754**, 214–234 (2016).
- [561] G. Aad (ATLAS), “Evidence for single top-quark production in the s -channel in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector using the Matrix Element Method”, *Phys. Lett. B* **756**, 228–246 (2016).
- [562] G. Aad (ATLAS), “Identification of boosted, hadronically decaying W bosons and comparisons with ATLAS data taken at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76**, 154 (2016).
- [563] G. Aad (ATLAS), “Identification of high transverse momentum top quarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **06**, 093 (2016).
- [564] G. Aad (ATLAS), “Measurement of $D^{*\pm}$, D^\pm and D_s^\pm meson production cross sections in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Nucl. Phys. B* **907**, 717–763 (2016).
- [565] G. Aad (ATLAS), “Measurement of W^\pm and Z -boson production cross sections in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **759**, 601–621 (2016).
- [566] G. Aad (ATLAS), “Measurement of event-shape observables in $Z \rightarrow \ell^+\ell^-$ events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC”, *Eur. Phys. J. C* **76**, 375 (2016).
- [567] G. Aad (ATLAS), “Measurement of fiducial differential cross sections of gluon-fusion production of Higgs bosons decaying to $WW \rightarrow e\nu\mu\nu$ with the ATLAS detector at $\sqrt{s} = 8$ TeV”, *JHEP* **08**, 104 (2016).
- [568] G. Aad (ATLAS), “Measurement of jet charge in dijet events from $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **93**, 052003 (2016).
- [569] G. Aad (ATLAS), “Measurement of the ZZ Production Cross Section in pp Collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **116**, 101801 (2016).
- [570] G. Aad (ATLAS), “Measurement of the angular coefficients in Z -boson events using electron and muon pairs from data taken at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **08**, 159 (2016).
- [571] G. Aad (ATLAS), “Measurement of the centrality dependence of the charged-particle pseudorapidity distribution in proton–lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, 199 (2016).
- [572] G. Aad (ATLAS), “Measurement of the charge asymmetry in highly boosted top-quark pair production in $\sqrt{s} = 8$ TeV pp collision data collected by the ATLAS experiment”, *Phys. Lett. B* **756**, 52–71 (2016).
- [573] G. Aad (ATLAS), “Measurement of the charge asymmetry in top-quark pair production in the lepton-plus-jets final state in pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, [Erratum: *Eur.Phys.J.C* 77, 564 (2017)], 87 (2016).
- [574] G. Aad (ATLAS), “Measurement of the charged-particle multiplicity inside jets from $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **76**, 322 (2016).
- [575] G. Aad (ATLAS), “Measurement of the correlations between the polar angles of leptons from top quark decays in the helicity basis at $\sqrt{s} = 7$ TeV using the ATLAS detector”, *Phys. Rev. D* **93**, 012002 (2016).
- [576] G. Aad (ATLAS), “Measurement of the CP-violating phase ϕ_s and the B_s^0 meson decay width difference with $B_s^0 \rightarrow J/\psi\phi$ decays in ATLAS”, *JHEP* **08**, 147 (2016).
- [577] G. Aad (ATLAS), “Measurement of the dependence of transverse energy production at large pseudorapidity on the hard-scattering kinematics of proton-proton collisions at $\sqrt{s} = 2.76$ TeV with ATLAS”, *Phys. Lett. B* **756**, 10–28 (2016).

- [578] G. Aad (ATLAS), “Measurement of the differential cross-section of highly boosted top quarks as a function of their transverse momentum in $\sqrt{s} = 8$ TeV proton-proton collisions using the ATLAS detector”, *Phys. Rev. D* **93**, 032009 (2016).
- [579] G. Aad (ATLAS), “Measurement of the differential cross-sections of prompt and non-prompt production of J/ψ and $\psi(2S)$ in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, 283 (2016).
- [580] G. Aad (ATLAS), “Measurement of the double-differential high-mass Drell-Yan cross section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **08**, 009 (2016).
- [581] G. Aad (ATLAS), “Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **08**, 005 (2016).
- [582] G. Aad (ATLAS), “Measurement of the production cross-section of a single top quark in association with a W boson at 8 TeV with the ATLAS experiment”, *JHEP* **01**, 064 (2016).
- [583] G. Aad (ATLAS), “Measurement of the transverse momentum and ϕ_{η}^* distributions of Drell-Yan lepton pairs in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **76**, 291 (2016).
- [584] G. Aad (ATLAS), “Measurement of total and differential W^+W^- production cross sections in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and limits on anomalous triple-gauge-boson couplings”, *JHEP* **09**, 029 (2016).
- [585] G. Aad (ATLAS), “Measurements of $W^{\pm}Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and limits on anomalous gauge boson self-couplings”, *Phys. Rev. D* **93**, 092004 (2016).
- [586] G. Aad (ATLAS), “Measurements of $Z\gamma$ and $Z\gamma\gamma$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **93**, 112002 (2016).
- [587] G. Aad (ATLAS), “Measurements of fiducial cross-sections for $t\bar{t}$ production with one or two additional b-jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **76**, 11 (2016).
- [588] G. Aad (ATLAS), “Measurements of four-lepton production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **753**, 552–572 (2016).
- [589] G. Aad (ATLAS), “Measurements of the charge asymmetry in top-quark pair production in the dilepton final state at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **94**, 032006 (2016).
- [590] G. Aad (ATLAS, CMS), “Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **08**, 045 (2016).
- [591] G. Aad (ATLAS), “Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $\sqrt{s} = 7$ and 8 TeV in the ATLAS experiment”, *Eur. Phys. J. C* **76**, 6 (2016).
- [592] G. Aad (ATLAS), “Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **76**, 538 (2016).
- [593] G. Aad (ATLAS), “Muon reconstruction performance of the ATLAS detector in proton-proton collision data at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **76**, 292 (2016).
- [594] G. Aad (ATLAS), “Observation of Long-Range Elliptic Azimuthal Anisotropies in $\sqrt{s} = 13$ and 2.76 TeV pp Collisions with the ATLAS Detector”, *Phys. Rev. Lett.* **116**, 172301 (2016).
- [595] G. Aad (ATLAS), “Performance of b -Jet Identification in the ATLAS Experiment”, *JINST* **11**, P04008 (2016).
- [596] G. Aad (ATLAS), “Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **76**, 581 (2016).
- [597] G. Aad (ATLAS), “Probing lepton flavour violation via neutrinoless $\tau \rightarrow 3\mu$ decays with the ATLAS detector”, *Eur. Phys. J. C* **76**, 232 (2016).
- [598] G. Aad (ATLAS), “Reconstruction of hadronic decay products of tau leptons with the ATLAS experiment”, *Eur. Phys. J. C* **76**, 295 (2016).
- [599] G. Aad (ATLAS), “Search for a high-mass Higgs boson decaying to a W boson pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **01**, 032 (2016).
- [600] G. Aad (ATLAS), “Search for an additional, heavy Higgs boson in the $H \rightarrow ZZ$ decay channel at $\sqrt{s} = 8$ TeV in pp collision data with the ATLAS detector”, *Eur. Phys. J. C* **76**, 45 (2016).
- [601] G. Aad (ATLAS), “Search for anomalous couplings in the Wtb vertex from the measurement of double differential angular decay rates of single top quarks produced in the t -channel with the ATLAS detector”, *JHEP* **04**, 023 (2016).

- [602] G. Aad (ATLAS), “Search for charged Higgs bosons in the $H^\pm \rightarrow tb$ decay channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, JHEP **03**, 127 (2016).
- [603] G. Aad (ATLAS), “Search for dark matter produced in association with a Higgs boson decaying to two bottom quarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. D **93**, 072007 (2016).
- [604] G. Aad (ATLAS), “Search for direct top squark pair production in final states with two tau leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 81 (2016).
- [605] G. Aad (ATLAS), “Search for flavour-changing neutral current top-quark decays to qZ in pp collision data collected with the ATLAS detector at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **76**, 12 (2016).
- [606] G. Aad (ATLAS), “Search for gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 565 (2016).
- [607] G. Aad (ATLAS), “Search for invisible decays of a Higgs boson using vector-boson fusion in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **01**, 172 (2016).
- [608] G. Aad (ATLAS), “Search for magnetic monopoles and stable particles with high electric charges in 8 TeV pp collisions with the ATLAS detector”, Phys. Rev. D **93**, 052009 (2016).
- [609] G. Aad (ATLAS), “Search for new phenomena in dijet mass and angular distributions from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. B **754**, 302–322 (2016).
- [610] G. Aad (ATLAS), “Search for new phenomena in events with at least three photons collected in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 210 (2016).
- [611] G. Aad (ATLAS), “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum with ATLAS using $\sqrt{s} = 13$ TeV proton-proton collisions”, Phys. Lett. B **757**, 334–355 (2016).
- [612] G. Aad (ATLAS), “Search for new phenomena with photon+jet events in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **03**, 041 (2016).
- [613] G. Aad (ATLAS), “Search for pair production of gluinos decaying via stop and sbottom in events with b -jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **94**, 032003 (2016).
- [614] G. Aad (ATLAS), “Search for single production of a vector-like quark via a heavy gluon in the $4b$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **758**, 249–268 (2016).
- [615] G. Aad (ATLAS), “Search for single production of vector-like quarks decaying into Wb in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 442 (2016).
- [616] G. Aad (ATLAS), “Search for strong gravity in multijet final states produced in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC”, JHEP **03**, 026 (2016).
- [617] G. Aad (ATLAS), “Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector”, Eur. Phys. J. C **76**, 259 (2016).
- [618] G. Aad (ATLAS), “Search for the electroweak production of supersymmetric particles in $\sqrt{s}=8$ TeV pp collisions with the ATLAS detector”, Phys. Rev. D **93**, 052002 (2016).
- [619] G. Aad (ATLAS), “Search for the production of single vector-like and excited quarks in the Wt final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **02**, 110 (2016).
- [620] G. Aad (ATLAS), “Search for the Standard Model Higgs boson decaying into $b\bar{b}$ produced in association with top quarks decaying hadronically in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **05**, 160 (2016).
- [621] G. Aad (ATLAS), “Search for the Standard Model Higgs boson produced in association with a vector boson and decaying into a tau pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. D **93**, 092005 (2016).
- [622] G. Aad (ATLAS), “Searches for scalar leptoquarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**, 5 (2016).
- [623] G. Aad (ATLAS), “Study of the $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays with the ATLAS detector”, Eur. Phys. J. C **76**, 4 (2016).
- [624] G. Aad (ATLAS), “Test of CP Invariance in vector-boson fusion production of the Higgs boson using the Optimal Observable method in the ditau decay channel with the ATLAS detector”, Eur. Phys. J. C **76**, 658 (2016).
- [625] G. Aad (ATLAS), “The performance of the jet trigger for the ATLAS detector during 2011 data taking”, Eur. Phys. J. C **76**, 526 (2016).

- [626] G. Aad (ATLAS), “Transverse momentum, rapidity, and centrality dependence of inclusive charged-particle production in $\sqrt{s_{NN}} = 5.02$ TeV $p + \text{Pb}$ collisions measured by the ATLAS experiment”, Phys. Lett. B **763**, 313–336 (2016).
- [627] M. Abolins (ATLAS TDAQ), “The ATLAS Data Acquisition and High Level Trigger system”, JINST **11**, P06008 (2016).
- [628] G. Aad (ATLAS), “ Z boson production in $p+\text{Pb}$ collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured with the ATLAS detector”, Phys. Rev. C **92**, 044915 (2015).
- [629] G. Aad (ATLAS), “A search for $t\bar{t}$ resonances using lepton-plus-jets events in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **08**, 148 (2015).
- [630] G. Aad (ATLAS), “A search for high-mass resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **07**, 157 (2015).
- [631] G. Aad (ATLAS), “Analysis of events with b -jets and a pair of leptons of the same charge in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **10**, 150 (2015).
- [632] G. Aad (ATLAS), “ATLAS Run 1 searches for direct pair production of third-generation squarks at the Large Hadron Collider”, Eur. Phys. J. C **75**, [Erratum: Eur.Phys.J.C 76, 153 (2016)], 510 (2015).
- [633] G. Aad (ATLAS), “Centrality and rapidity dependence of inclusive jet production in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector”, Phys. Lett. B **748**, 392–413 (2015).
- [634] G. Aad (ATLAS, CMS), “Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments”, Phys. Rev. Lett. **114**, 191803 (2015).
- [635] G. Aad (ATLAS), “Constraints on new phenomena via Higgs boson couplings and invisible decays with the ATLAS detector”, JHEP **11**, 206 (2015).
- [636] G. Aad (ATLAS), “Constraints on the off-shell Higgs boson signal strength in the high-mass ZZ and WW final states with the ATLAS detector”, Eur. Phys. J. C **75**, 335 (2015).
- [637] G. Aad (ATLAS), “Determination of spin and parity of the Higgs boson in the $WW^* \rightarrow e\nu\mu\nu$ decay channel with the ATLAS detector”, Eur. Phys. J. C **75**, 231 (2015).
- [638] G. Aad (ATLAS), “Determination of the ratio of b -quark fragmentation fractions f_s/f_d in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **115**, 262001 (2015).
- [639] G. Aad (ATLAS), “Determination of the top-quark pole mass using $t\bar{t} + 1$ -jet events collected with the ATLAS experiment in 7 TeV pp collisions”, JHEP **10**, 121 (2015).
- [640] G. Aad (ATLAS), “Differential top-antitop cross-section measurements as a function of observables constructed from final-state particles using pp collisions at $\sqrt{s} = 7$ TeV in the ATLAS detector”, JHEP **06**, 100 (2015).
- [641] G. Aad (ATLAS), “Evidence for the Higgs-boson Yukawa coupling to tau leptons with the ATLAS detector”, JHEP **04**, 117 (2015).
- [642] G. Aad (ATLAS), “Evidence of $W\gamma\gamma$ Production in pp Collisions at $s=8$ TeV and Limits on Anomalous Quartic Gauge Couplings with the ATLAS Detector”, Phys. Rev. Lett. **115**, 031802 (2015).
- [643] G. Aad (ATLAS), “Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at $\sqrt{s}=8$ TeV”, Eur. Phys. J. C **75**, 303 (2015).
- [644] G. Aad (ATLAS), “Jet energy measurement and its systematic uncertainty in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **75**, 17 (2015).
- [645] G. Aad (ATLAS), “Measurement of charged-particle spectra in $\text{Pb}+\text{Pb}$ collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC”, JHEP **09**, 050 (2015).
- [646] G. Aad (ATLAS), “Measurement of colour flow with the jet pull angle in $t\bar{t}$ events using the ATLAS detector at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **750**, 475–493 (2015).
- [647] G. Aad (ATLAS), “Measurement of differential J/ψ production cross sections and forward-backward ratios in $p + \text{Pb}$ collisions with the ATLAS detector”, Phys. Rev. C **92**, 034904 (2015).
- [648] G. Aad (ATLAS), “Measurement of exclusive $\gamma\gamma \rightarrow \ell^+\ell^-$ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **749**, 242–261 (2015).
- [649] G. Aad (ATLAS), “Measurement of four-jet differential cross sections in $\sqrt{s} = 8$ TeV proton-proton collisions using the ATLAS detector”, JHEP **12**, 105 (2015).
- [650] G. Aad (ATLAS), “Measurement of Spin Correlation in Top-Antitop Quark Events and Search for Top Squark Pair Production in pp Collisions at $\sqrt{s} = 8$ TeV Using the ATLAS Detector”, Phys. Rev. Lett. **114**, 142001 (2015).
- [651] G. Aad (ATLAS), “Measurement of the $t\bar{t}$ production cross-section as a function of jet multiplicity and jet transverse momentum in 7 TeV proton-proton collisions with the ATLAS detector”, JHEP **01**, 020 (2015).

- [652] G. Aad (ATLAS), “Measurement of the $t\bar{t}W$ and $t\bar{t}Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **11**, 172 (2015).
- [653] G. Aad (ATLAS), “Measurement of the $WW + WZ$ cross section and limits on anomalous triple gauge couplings using final states with one lepton, missing transverse momentum, and two jets with the ATLAS detector at $\sqrt{s} = 7$ TeV”, JHEP **01**, 049 (2015).
- [654] G. Aad (ATLAS), “Measurement of the branching ratio $\Gamma(\Lambda_b^0 \rightarrow \psi(2S)\Lambda^0)/\Gamma(\Lambda_b^0 \rightarrow J/\psi\Lambda^0)$ with the ATLAS detector”, Phys. Lett. B **751**, 63–80 (2015).
- [655] G. Aad (ATLAS), “Measurement of the charge asymmetry in dileptonic decays of top quark pairs in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **05**, 061 (2015).
- [656] G. Aad (ATLAS), “Measurement of the correlation between flow harmonics of different order in lead-lead collisions at $\sqrt{s_{NN}}=2.76$ TeV with the ATLAS detector”, Phys. Rev. C **92**, 034903 (2015).
- [657] G. Aad (ATLAS), “Measurement of the forward-backward asymmetry of electron and muon pair-production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **09**, 049 (2015).
- [658] G. Aad (ATLAS), “Measurement of the inclusive jet cross-section in proton-proton collisions at $\sqrt{s} = 7$ TeV using 4.5 fb⁻¹ of data with the ATLAS detector”, JHEP **02**, [Erratum: JHEP 09, 141 (2015)], 153 (2015).
- [659] G. Aad (ATLAS), “Measurement of the production and lepton charge asymmetry of W bosons in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Eur. Phys. J. C **75**, 23 (2015).
- [660] G. Aad (ATLAS), “Measurement of the production of neighbouring jets in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. B **751**, 376–395 (2015).
- [661] G. Aad (ATLAS), “Measurement of the top pair production cross section in 8 TeV proton-proton collisions using kinematic information in the lepton+jets final state with ATLAS”, Phys. Rev. D **91**, 112013 (2015).
- [662] G. Aad (ATLAS), “Measurement of the top quark mass in the $t\bar{t} \rightarrow$ lepton+jets and $t\bar{t} \rightarrow$ dilepton channels using $\sqrt{s} = 7$ TeV ATLAS data”, Eur. Phys. J. C **75**, 330 (2015).
- [663] G. Aad (ATLAS), “Measurement of the top-quark mass in the fully hadronic decay channel from ATLAS data at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **75**, 158 (2015).
- [664] G. Aad (ATLAS), “Measurement of the transverse polarization of Λ and $\bar{\Lambda}$ hyperons produced in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. D **91**, 032004 (2015).
- [665] G. Aad (ATLAS), “Measurement of three-jet production cross-sections in pp collisions at 7 TeV centre-of-mass energy using the ATLAS detector”, Eur. Phys. J. C **75**, 228 (2015).
- [666] G. Aad (ATLAS), “Measurement of transverse energy-energy correlations in multi-jet events in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector and determination of the strong coupling constant $\alpha_s(m_Z)$ ”, Phys. Lett. B **750**, 427–447 (2015).
- [667] G. Aad (ATLAS), “Measurements of Higgs boson production and couplings in the four-lepton channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector”, Phys. Rev. D **91**, 012006 (2015).
- [668] G. Aad (ATLAS), “Measurements of the Nuclear Modification Factor for Jets in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **114**, 072302 (2015).
- [669] G. Aad (ATLAS), “Measurements of the top quark branching ratios into channels with leptons and quarks with the ATLAS detector”, Phys. Rev. D **92**, 072005 (2015).
- [670] G. Aad (ATLAS), “Measurements of the Total and Differential Higgs Boson Production Cross Sections Combining the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ Decay Channels at $\sqrt{s}=8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **115**, 091801 (2015).
- [671] G. Aad (ATLAS), “Measurements of the W production cross sections in association with jets with the ATLAS detector”, Eur. Phys. J. C **75**, 82 (2015).
- [672] G. Aad (ATLAS), “Modelling $Z \rightarrow \tau\tau$ processes in ATLAS with τ -embedded $Z \rightarrow \mu\mu$ data”, JINST **10**, P09018 (2015).
- [673] G. Aad (ATLAS), “Observation and measurement of Higgs boson decays to WW^* with the ATLAS detector”, Phys. Rev. D **92**, 012006 (2015).
- [674] G. Aad (ATLAS), “Observation and measurements of the production of prompt and non-prompt J/ψ mesons in association with a Z boson in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **75**, 229 (2015).
- [675] G. Aad (ATLAS), “Observation of top-quark pair production in association with a photon and measurement of the $t\bar{t}\gamma$ production cross section in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. D **91**, 072007 (2015).

- [676] G. Aad (ATLAS), “Performance of the ATLAS muon trigger in pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **75**, 120 (2015).
- [677] G. Aad (ATLAS), “Search for $H \rightarrow \gamma\gamma$ produced in association with top quarks and constraints on the Yukawa coupling between the top quark and the Higgs boson using data taken at 7 TeV and 8 TeV with the ATLAS detector”, Phys. Lett. B **740**, 222–242 (2015).
- [678] G. Aad (ATLAS), “Search for s -channel single top-quark production in proton–proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B **740**, 118–136 (2015).
- [679] G. Aad (ATLAS), “Search for $W' \rightarrow tb \rightarrow qqbb$ decays in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **75**, 165 (2015).
- [680] G. Aad (ATLAS), “Search for $W' \rightarrow t\bar{b}$ in the lepton plus jets final state in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B **743**, 235–255 (2015).
- [681] G. Aad (ATLAS), “Search for a Charged Higgs Boson Produced in the Vector-Boson Fusion Mode with Decay $H^\pm \rightarrow W^\pm Z$ using pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Experiment”, Phys. Rev. Lett. **114**, 231801 (2015).
- [682] G. Aad (ATLAS), “Search for a CP-odd Higgs boson decaying to Zh in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B **744**, 163–183 (2015).
- [683] G. Aad (ATLAS), “Search for a Heavy Neutral Particle Decaying to $e\mu$, $e\tau$, or $\mu\tau$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **115**, 031801 (2015).
- [684] G. Aad (ATLAS), “Search for a new resonance decaying to a W or Z boson and a Higgs boson in the $\ell\ell/\ell\nu/\nu\nu + b\bar{b}$ final states with the ATLAS detector”, Eur. Phys. J. C **75**, 263 (2015).
- [685] G. Aad (ATLAS), “Search for anomalous production of prompt same-sign lepton pairs and pair-produced doubly charged Higgs bosons with $\sqrt{s} = 8$ TeV pp collisions using the ATLAS detector”, JHEP **03**, 041 (2015).
- [686] G. Aad (ATLAS), “Search for charged Higgs bosons decaying via $H^\pm \rightarrow \tau^\pm\nu$ in fully hadronic final states using pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **03**, 088 (2015).
- [687] G. Aad (ATLAS), “Search for dark matter in events with heavy quarks and missing transverse momentum in pp collisions with the ATLAS detector”, Eur. Phys. J. C **75**, 92 (2015).
- [688] G. Aad (ATLAS), “Search for Dark Matter in Events with Missing Transverse Momentum and a Higgs Boson Decaying to Two Photons in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **115**, 131801 (2015).
- [689] G. Aad (ATLAS), “Search for direct pair production of a chargino and a neutralino decaying to the 125 GeV Higgs boson in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **75**, 208 (2015).
- [690] G. Aad (ATLAS), “Search for flavour-changing neutral current top quark decays $t \rightarrow Hq$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **12**, 061 (2015).
- [691] G. Aad (ATLAS), “Search for heavy lepton resonances decaying to a Z boson and a lepton in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 108 (2015).
- [692] G. Aad (ATLAS), “Search for heavy long-lived multi-charged particles in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Eur. Phys. J. C **75**, 362 (2015).
- [693] G. Aad (ATLAS), “Search for heavy Majorana neutrinos with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **07**, 162 (2015).
- [694] G. Aad (ATLAS), “Search for Higgs and Z Boson Decays to $J/\psi\gamma$ and $\Upsilon(nS)\gamma$ with the ATLAS Detector”, Phys. Rev. Lett. **114**, 121801 (2015).
- [695] G. Aad (ATLAS), “Search For Higgs Boson Pair Production in the $\gamma\gamma b\bar{b}$ Final State using pp Collision Data at $\sqrt{s} = 8$ TeV from the ATLAS Detector”, Phys. Rev. Lett. **114**, 081802 (2015).
- [696] G. Aad (ATLAS), “Search for Higgs boson pair production in the $b\bar{b}b\bar{b}$ final state from pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **75**, 412 (2015).
- [697] G. Aad (ATLAS), “Search for Higgs bosons decaying to aa in the $\mu\mu\tau\tau$ final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS experiment”, Phys. Rev. D **92**, 052002 (2015).
- [698] G. Aad (ATLAS), “Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **12**, 055 (2015).
- [699] G. Aad (ATLAS), “Search for high-mass diphoton resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. D **92**, 032004 (2015).
- [700] G. Aad (ATLAS), “Search for invisible decays of the Higgs boson produced in association with a hadronically decaying vector boson in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. C **75**, 337 (2015).

- [701] G. Aad (ATLAS), “Search for invisible particles produced in association with single-top-quarks in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **75**, 79 (2015).
- [702] G. Aad (ATLAS), “Search for lepton-flavour-violating $H \rightarrow \mu\tau$ decays of the Higgs boson with the ATLAS detector”, *JHEP* **11**, 211 (2015).
- [703] G. Aad (ATLAS), “Search for long-lived, weakly interacting particles that decay to displaced hadronic jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **92**, 012010 (2015).
- [704] G. Aad (ATLAS), “Search for low-scale gravity signatures in multi-jet final states with the ATLAS detector at $\sqrt{s} = 8$ TeV”, *JHEP* **07**, 032 (2015).
- [705] G. Aad (ATLAS), “Search for massive supersymmetric particles decaying to many jets using the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. D* **91**, [Erratum: *Phys.Rev.D* 93, 039901 (2016)], 112016 (2015).
- [706] G. Aad (ATLAS), “Search for massive, long-lived particles using multitrack displaced vertices or displaced lepton pairs in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **92**, 072004 (2015).
- [707] G. Aad (ATLAS), “Search for metastable heavy charged particles with large ionisation energy loss in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS experiment”, *Eur. Phys. J. C* **75**, 407 (2015).
- [708] G. Aad (ATLAS), “Search for new light gauge bosons in Higgs boson decays to four-lepton final states in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC”, *Phys. Rev. D* **92**, 092001 (2015).
- [709] G. Aad (ATLAS), “Search for New Phenomena in Dijet Angular Distributions in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV Measured with the ATLAS Detector”, *Phys. Rev. Lett.* **114**, 221802 (2015).
- [710] G. Aad (ATLAS), “Search for new phenomena in events with a photon and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **91**, [Erratum: *Phys.Rev.D* 92, 059903 (2015)], 012008 (2015).
- [711] G. Aad (ATLAS), “Search for new phenomena in events with three or more charged leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **08**, 138 (2015).
- [712] G. Aad (ATLAS), “Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **75**, [Erratum: *Eur.Phys.J.C* 75, 408 (2015)], 299 (2015).
- [713] G. Aad (ATLAS), “Search for new phenomena in the dijet mass distribution using $p - p$ collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **91**, 052007 (2015).
- [714] G. Aad (ATLAS), “Search for pair production of a new heavy quark that decays into a W boson and a light quark in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **92**, 112007 (2015).
- [715] G. Aad (ATLAS), “Search for pair-produced long-lived neutral particles decaying in the ATLAS hadronic calorimeter in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Lett. B* **743**, 15–34 (2015).
- [716] G. Aad (ATLAS), “Search for photonic signatures of gauge-mediated supersymmetry in 8 TeV pp collisions with the ATLAS detector”, *Phys. Rev. D* **92**, 072001 (2015).
- [717] G. Aad (ATLAS), “Search for production of WW/WZ resonances decaying to a lepton, neutrino and jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **75**, [Erratum: *Eur.Phys.J.C* 75, 370 (2015)], 209 (2015).
- [718] G. Aad (ATLAS), “Search for production of vector-like quark pairs and of four top quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **08**, 105 (2015).
- [719] G. Aad (ATLAS), “Search for resonant diboson production in the $\ell\ell q\bar{q}$ final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **75**, 69 (2015).
- [720] G. Aad (ATLAS), “Search for Scalar Charm Quark Pair Production in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **114**, 161801 (2015).
- [721] G. Aad (ATLAS), “Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **04**, 116 (2015).
- [722] G. Aad (ATLAS), “Search for supersymmetry in events containing a same-flavour opposite-sign dilepton pair, jets, and large missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **75**, [Erratum: *Eur.Phys.J.C* 75, 463 (2015)], 318 (2015).
- [723] G. Aad (ATLAS), “Search for the $b\bar{b}$ decay of the Standard Model Higgs boson in associated $(W/Z)H$ production with the ATLAS detector”, *JHEP* **01**, 069 (2015).
- [724] G. Aad (ATLAS), “Search for the X_b and other hidden-beauty states in the $\pi^+\pi^-\Upsilon(1S)$ channel at ATLAS”, *Phys. Lett. B* **740**, 199–217 (2015).

- [725] G. Aad (ATLAS), “Search for the associated production of the Higgs boson with a top quark pair in multilepton final states with the ATLAS detector”, *Phys. Lett. B* **749**, 519–541 (2015).
- [726] G. Aad (ATLAS), “Search for the Standard Model Higgs boson produced in association with top quarks and decaying into $b\bar{b}$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **75**, 349 (2015).
- [727] G. Aad (ATLAS), “Search for type-III Seesaw heavy leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, *Phys. Rev. D* **92**, 032001 (2015).
- [728] G. Aad (ATLAS), “Search for vector-like B quarks in events with one isolated lepton, missing transverse momentum and jets at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **91**, 112011 (2015).
- [729] G. Aad (ATLAS), “Searches for heavy long-lived charged particles with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **01**, 068 (2015).
- [730] G. Aad (ATLAS), “Searches for Higgs boson pair production in the $hh \rightarrow bb\tau\tau, \gamma\gamma WW^*, \gamma\gamma bb, bbbb$ channels with the ATLAS detector”, *Phys. Rev. D* **92**, 092004 (2015).
- [731] G. Aad (ATLAS), “Simultaneous measurements of the $t\bar{t}$, W^+W^- , and $Z/\gamma^* \rightarrow \tau\tau$ production cross-sections in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Rev. D* **91**, 052005 (2015).
- [732] G. Aad (ATLAS), “Study of (W/Z)H production and Higgs boson couplings using $H \rightarrow WW^*$ decays with the ATLAS detector”, *JHEP* **08**, 137 (2015).
- [733] G. Aad (ATLAS), “Study of the spin and parity of the Higgs boson in diboson decays with the ATLAS detector”, *Eur. Phys. J. C* **75**, [Erratum: *Eur.Phys.J.C* 76, 152 (2016)], 476 (2015).
- [734] G. Aad (ATLAS), “Summary of the ATLAS experiment’s sensitivity to supersymmetry after LHC Run 1 — interpreted in the phenomenological MSSM”, *JHEP* **10**, 134 (2015).
- [735] G. Aad (ATLAS), “Summary of the searches for squarks and gluinos using $\sqrt{s} = 8$ TeV pp collisions with the ATLAS experiment at the LHC”, *JHEP* **10**, 054 (2015).
- [736] G. Aad (ATLAS), “Two-particle Bose–Einstein correlations in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector”, *Eur. Phys. J. C* **75**, 466 (2015).
- [737] G. Volpi, “The ATLAS fast tracker processor design”, *PoS VERTEX2015*, 040 (2015).
- [738] G. Aad (ATLAS), “A measurement of the ratio of the production cross sections for W and Z bosons in association with jets with the ATLAS detector”, *Eur. Phys. J. C* **74**, 3168 (2014).
- [739] G. Aad (ATLAS), “A neural network clustering algorithm for the ATLAS silicon pixel detector”, *JINST* **9**, P09009 (2014).
- [740] G. Aad (ATLAS), “Comprehensive measurements of t -channel single top-quark production cross sections at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Rev. D* **90**, 112006 (2014).
- [741] G. Aad (ATLAS), “Electron and photon energy calibration with the ATLAS detector using LHC Run 1 data”, *Eur. Phys. J. C* **74**, 3071 (2014).
- [742] G. Aad (ATLAS), “Electron reconstruction and identification efficiency measurements with the ATLAS detector using the 2011 LHC proton-proton collision data”, *Eur. Phys. J. C* **74**, 2941 (2014).
- [743] G. Aad (ATLAS), “Evidence for Electroweak Production of $W^\pm W^\pm jj$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **113**, 141803 (2014).
- [744] G. Aad (ATLAS), “Fiducial and differential cross sections of Higgs boson production measured in the four-lepton decay channel in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector”, *Phys. Lett. B* **738**, 234–253 (2014).
- [745] G. Aad (ATLAS), “Flavor tagged time-dependent angular analysis of the $B_s \rightarrow J/\psi\phi$ decay and extraction of $\Delta\Gamma_s$ and the weak phase ϕ_s in ATLAS”, *Phys. Rev. D* **90**, 052007 (2014).
- [746] G. Aad (ATLAS), “Light-quark and gluon jet discrimination in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **74**, 3023 (2014).
- [747] G. Aad (ATLAS), “Measurement of χ_{e1} and χ_{e2} production with $\sqrt{s} = 7$ TeV pp collisions at ATLAS”, *JHEP* **07**, 154 (2014).
- [748] G. Aad (ATLAS), “Measurement of differential production cross-sections for a Z boson in association with b -jets in 7 TeV proton-proton collisions with the ATLAS detector”, *JHEP* **10**, 141 (2014).
- [749] G. Aad (ATLAS), “Measurement of dijet cross sections in pp collisions at 7 TeV centre-of-mass energy using the ATLAS detector”, *JHEP* **05**, 059 (2014).
- [750] G. Aad (ATLAS), “Measurement of distributions sensitive to the underlying event in inclusive Z -boson production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **74**, 3195 (2014).

- [751] G. Aad (ATLAS), “Measurement of event-plane correlations in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector”, Phys. Rev. C **90**, 024905 (2014).
- [752] G. Aad (ATLAS), “Measurement of flow harmonics with multi-particle cumulants in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Eur. Phys. J. C **74**, 3157 (2014).
- [753] G. Aad (ATLAS), “Measurement of Higgs boson production in the diphoton decay channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector”, Phys. Rev. D **90**, 112015 (2014).
- [754] G. Aad (ATLAS), “Measurement of inclusive jet charged-particle fragmentation functions in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. B **739**, 320–342 (2014).
- [755] G. Aad (ATLAS), “Measurement of long-range pseudorapidity correlations and azimuthal harmonics in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector”, Phys. Rev. C **90**, 044906 (2014).
- [756] G. Aad (ATLAS), “Measurement of the $t\bar{t}$ production cross-section using $e\mu$ events with b-tagged jets in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS detector”, Eur. Phys. J. C **74**, [Addendum: Eur.Phys.J.C 76, 642 (2016)], 3109 (2014).
- [757] G. Aad (ATLAS), “Measurement of the Z/γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **09**, 145 (2014).
- [758] G. Aad (ATLAS), “Measurement of the centrality and pseudorapidity dependence of the integrated elliptic flow in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Eur. Phys. J. C **74**, 2982 (2014).
- [759] G. Aad (ATLAS), “Measurement of the cross section of high transverse momentum $Z \rightarrow b\bar{b}$ production in proton–proton collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS Detector”, Phys. Lett. B **738**, 25–43 (2014).
- [760] G. Aad (ATLAS), “Measurement of the cross-section of high transverse momentum vector bosons reconstructed as single jets and studies of jet substructure in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, New J. Phys. **16**, 113013 (2014).
- [761] G. Aad (ATLAS), “Measurement of the electroweak production of dijets in association with a Z-boson and distributions sensitive to vector boson fusion in proton-proton collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, JHEP **04**, 031 (2014).
- [762] G. Aad (ATLAS), “Measurement of the Higgs boson mass from the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ channels with the ATLAS detector using 25fb^{-1} of pp collision data”, Phys. Rev. D **90**, 052004 (2014).
- [763] G. Aad (ATLAS), “Measurement of the inclusive isolated prompt photons cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector using 4.6fb^{-1} ”, Phys. Rev. D **89**, 052004 (2014).
- [764] G. Aad (ATLAS), “Measurement of the low-mass Drell-Yan differential cross section at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **06**, 112 (2014).
- [765] G. Aad (ATLAS), “Measurement of the mass difference between top and anti-top quarks in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Lett. B **728**, 363–379 (2014).
- [766] G. Aad (ATLAS), “Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton–proton collision data”, Eur. Phys. J. C **74**, 3130 (2014).
- [767] G. Aad (ATLAS), “Measurement of the parity-violating asymmetry parameter α_b and the helicity amplitudes for the decay $\Lambda_b^0 \rightarrow J/\psi + \Lambda^0$ with the ATLAS detector”, Phys. Rev. D **89**, 092009 (2014).
- [768] G. Aad (ATLAS), “Measurement of the production cross section of prompt J/ψ mesons in association with a W^\pm boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **04**, 172 (2014).
- [769] G. Aad (ATLAS), “Measurement of the production cross-section of $\psi(2S) \rightarrow J/\psi(\rightarrow \mu^+\mu^-)\pi^+\pi^-$ in pp collisions at $\sqrt{s} = 7$ TeV at ATLAS”, JHEP **09**, 079 (2014).
- [770] G. Aad (ATLAS), “Measurement of the production of a W boson in association with a charm quark in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **05**, 068 (2014).
- [771] G. Aad (ATLAS), “Measurement of the top quark pair production charge asymmetry in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **02**, 107 (2014).
- [772] G. Aad (ATLAS), “Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. B **889**, 486–548 (2014).
- [773] G. Aad (ATLAS), “Measurement of the underlying event in jet events from 7 TeV proton-proton collisions with the ATLAS detector”, Eur. Phys. J. C **74**, 2965 (2014).
- [774] G. Aad (ATLAS), “Measurements of fiducial and differential cross sections for Higgs boson production in the diphoton decay channel at $\sqrt{s} = 8$ TeV with ATLAS”, JHEP **09**, 112 (2014).
- [775] G. Aad (ATLAS), “Measurements of Four-Lepton Production at the Z Resonance in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS”, Phys. Rev. Lett. **112**, 231806 (2014).

- [776] G. Aad (ATLAS), “Measurements of jet vetoes and azimuthal decorrelations in dijet events produced in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **74**, 3117 (2014).
- [777] G. Aad (ATLAS), “Measurements of normalized differential cross sections for $t\bar{t}$ production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, *Phys. Rev. D* **90**, 072004 (2014).
- [778] G. Aad (ATLAS), “Measurements of spin correlation in top-antitop quark events from proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, *Phys. Rev. D* **90**, 112016 (2014).
- [779] G. Aad (ATLAS), “Monitoring and data quality assessment of the ATLAS liquid argon calorimeter”, *JINST* **9**, P07024 (2014).
- [780] G. Aad (ATLAS), “Muon reconstruction efficiency and momentum resolution of the ATLAS experiment in proton-proton collisions at $\sqrt{s} = 7$ TeV in 2010”, *Eur. Phys. J. C* **74**, 3034 (2014).
- [781] G. Aad (ATLAS), “Observation of an Excited B_c^\pm Meson State with the ATLAS Detector”, *Phys. Rev. Lett.* **113**, 212004 (2014).
- [782] G. Aad (ATLAS), “Operation and performance of the ATLAS semiconductor tracker”, *JINST* **9**, P08009 (2014).
- [783] G. Aad (ATLAS), “Search for a multi-Higgs-boson cascade in $W^+W^b\bar{b}$ events with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. D* **89**, 032002 (2014).
- [784] G. Aad (ATLAS), “Search for contact interactions and large extra dimensions in the dilepton channel using proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **74**, 3134 (2014).
- [785] G. Aad (ATLAS), “Search for dark matter in events with a hadronically decaying W or Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **112**, 041802 (2014).
- [786] G. Aad (ATLAS), “Search for dark matter in events with a Z boson and missing transverse momentum in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector”, *Phys. Rev. D* **90**, 012004 (2014).
- [787] G. Aad (ATLAS), “Search for direct pair production of the top squark in all-hadronic final states in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **09**, 015 (2014).
- [788] G. Aad (ATLAS), “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *JHEP* **04**, 169 (2014).
- [789] G. Aad (ATLAS), “Search for direct production of charginos, neutralinos and sleptons in final states with two leptons and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **05**, 071 (2014).
- [790] G. Aad (ATLAS), “Search for direct top squark pair production in events with a Z boson, b-jets and missing transverse momentum in $\sqrt{s}=8$ TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **74**, 2883 (2014).
- [791] G. Aad (ATLAS), “Search for direct top-squark pair production in final states with two leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **06**, 124 (2014).
- [792] G. Aad (ATLAS), “Search for Higgs boson decays to a photon and a Z boson in pp collisions at $\sqrt{s}=7$ and 8 TeV with the ATLAS detector”, *Phys. Lett. B* **732**, 8–27 (2014).
- [793] G. Aad (ATLAS), “Search for high-mass dilepton resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev. D* **90**, 052005 (2014).
- [794] G. Aad (ATLAS), “Search for Invisible Decays of a Higgs Boson Produced in Association with a Z Boson in ATLAS”, *Phys. Rev. Lett.* **112**, 201802 (2014).
- [795] G. Aad (ATLAS), “Search for long-lived neutral particles decaying into lepton jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **11**, 088 (2014).
- [796] G. Aad (ATLAS), “Search for microscopic black holes and string balls in final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 8$ TeV”, *JHEP* **08**, 103 (2014).
- [797] G. Aad (ATLAS), “Search for neutral Higgs bosons of the minimal supersymmetric standard model in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **11**, 056 (2014).
- [798] G. Aad (ATLAS), “Search for new particles in events with one lepton and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *JHEP* **09**, 037 (2014).
- [799] G. Aad (ATLAS), “Search for new phenomena in photon+jet events collected in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **728**, 562–578 (2014).
- [800] G. Aad (ATLAS), “Search for new resonances in $W\gamma$ and $Z\gamma$ final states in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **738**, 428–447 (2014).

- [801] G. Aad (ATLAS), “Search for nonpointing and delayed photons in the diphoton and missing transverse momentum final state in 8 TeV pp collisions at the LHC using the ATLAS detector”, Phys. Rev. D **90**, 112005 (2014).
- [802] G. Aad (ATLAS), “Search for pair and single production of new heavy quarks that decay to a Z boson and a third-generation quark in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **11**, 104 (2014).
- [803] G. Aad (ATLAS), “Search for pair-produced third-generation squarks decaying via charm quarks or in compressed supersymmetric scenarios in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. D **90**, 052008 (2014).
- [804] G. Aad (ATLAS), “Search for Quantum Black Hole Production in High-Invariant-Mass Lepton+Jet Final States Using pp Collisions at $\sqrt{s} = 8$ TeV and the ATLAS Detector”, Phys. Rev. Lett. **112**, 091804 (2014).
- [805] G. Aad (ATLAS), “Search for Scalar Diphoton Resonances in the Mass Range 65 – 600 GeV with the ATLAS Detector in pp Collision Data at $\sqrt{s} = 8$ TeV”, Phys. Rev. Lett. **113**, 171801 (2014).
- [806] G. Aad (ATLAS), “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton–proton collision data”, JHEP **09**, 176 (2014).
- [807] G. Aad (ATLAS), “Search for strong production of supersymmetric particles in final states with missing transverse momentum and at least three b -jets at $\sqrt{s} = 8$ TeV proton-proton collisions with the ATLAS detector”, JHEP **10**, 024 (2014).
- [808] G. Aad (ATLAS), “Search for supersymmetry at $\sqrt{s} = 8$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector”, JHEP **06**, 035 (2014).
- [809] G. Aad (ATLAS), “Search for supersymmetry in events with four or more leptons in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Phys. Rev. D **90**, 052001 (2014).
- [810] G. Aad (ATLAS), “Search for supersymmetry in events with large missing transverse momentum, jets, and at least one tau lepton in 20 fb⁻¹ of $\sqrt{s} = 8$ TeV proton-proton collision data with the ATLAS detector”, JHEP **09**, 103 (2014).
- [811] G. Aad (ATLAS), “Search for the direct production of charginos, neutralinos and staus in final states with at least two hadronically decaying taus and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **10**, 096 (2014).
- [812] G. Aad (ATLAS), “Search for the lepton flavor violating decay $Z \rightarrow e\mu$ in pp collisions at \sqrt{s} TeV with the ATLAS detector”, Phys. Rev. D **90**, 072010 (2014).
- [813] G. Aad (ATLAS), “Search for the Standard Model Higgs boson decay to $\mu^+\mu^-$ with the ATLAS detector”, Phys. Lett. B **738**, 68–86 (2014).
- [814] G. Aad (ATLAS), “Search for top quark decays $t \rightarrow qH$ with $H \rightarrow \gamma\gamma$ using the ATLAS detector”, JHEP **06**, 008 (2014).
- [815] G. Aad (ATLAS), “Search for top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, JHEP **11**, 118 (2014).
- [816] G. Aad (ATLAS), “Search for WZ resonances in the fully leptonic channel using pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B **737**, 223–243 (2014).
- [817] G. Aad (ATLAS), “Standalone vertex finding in the ATLAS muon spectrometer”, JINST **9**, P02001 (2014).
- [818] G. Aad (ATLAS), “Study of heavy-flavor quarks produced in association with top-quark pairs at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. D **89**, 072012 (2014).
- [819] G. Aad (ATLAS), “The differential production cross section of the ϕ (1020) meson in $\sqrt{s} = 7$ TeV pp collisions measured with the ATLAS detector”, Eur. Phys. J. C **74**, 2895 (2014).
- [820] G. Aad (ATLAS), “A search for high-mass resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **719**, 242–260 (2013).
- [821] G. Aad (ATLAS), “A search for prompt lepton-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **719**, 299–317 (2013).
- [822] G. Aad (ATLAS), “ATLAS search for new phenomena in dijet mass and angular distributions using pp collisions at $\sqrt{s} = 7$ TeV”, JHEP **01**, 029 (2013).
- [823] G. Aad (ATLAS), “Characterisation and mitigation of beam-induced backgrounds observed in the ATLAS detector during the 2011 proton-proton run”, JINST **8**, P07004 (2013).

- [824] G. Aad (ATLAS), “Dynamics of isolated-photon plus jet production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. B **875**, 483–535 (2013).
- [825] G. Aad (ATLAS), “Evidence for the spin-0 nature of the Higgs boson using ATLAS data”, Phys. Lett. B **726**, 120–144 (2013).
- [826] G. Aad (ATLAS), “Improved luminosity determination in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector at the LHC”, Eur. Phys. J. C **73**, 2518 (2013).
- [827] G. Aad (ATLAS), “Jet energy measurement with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **73**, 2304 (2013).
- [828] G. Aad (ATLAS), “Jet energy resolution in proton-proton collisions at $\sqrt{s} = 7$ TeV recorded in 2010 with the ATLAS detector”, Eur. Phys. J. C **73**, 2306 (2013).
- [829] G. Aad (ATLAS), “Measurement of W^+W^- production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector and limits on anomalous WWZ and $WW\gamma$ couplings”, Phys. Rev. D **87**, [Erratum: Phys.Rev.D 88, 079906 (2013)], 112001 (2013).
- [830] G. Aad (ATLAS), “Measurement of Z boson Production in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **110**, 022301 (2013).
- [831] G. Aad (ATLAS), “Measurement of ZZ production in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous ZZZ and $ZZ\gamma$ couplings with the ATLAS detector”, JHEP **03**, 128 (2013).
- [832] G. Aad (ATLAS), “Measurement of angular correlations in Drell-Yan lepton pairs to probe Z/gamma* boson transverse momentum at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. B **720**, 32–51 (2013).
- [833] G. Aad (ATLAS), “Measurement of charged-particle event shape variables in $\sqrt{s} = 7$ TeV proton-proton interactions with the ATLAS detector”, Phys. Rev. D **88**, 032004 (2013).
- [834] G. Aad (ATLAS), “Measurement of hard double-parton interactions in $W(\rightarrow l\nu) + 2$ jet events at $\sqrt{s}=7$ TeV with the ATLAS detector”, New J. Phys. **15**, 033038 (2013).
- [835] G. Aad (ATLAS), “Measurement of isolated-photon pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **01**, 086 (2013).
- [836] G. Aad (ATLAS), “Measurement of jet shapes in top-quark pair events at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. C **73**, 2676 (2013).
- [837] G. Aad (ATLAS), “Measurement of kT splitting scales in $W \rightarrow l\nu$ events at $\sqrt{s}=7$ TeV with the ATLAS detector”, Eur. Phys. J. C **73**, 2432 (2013).
- [838] G. Aad (ATLAS), “Measurement of the Λ_b^0 lifetime and mass in the ATLAS experiment”, Phys. Rev. D **87**, 032002 (2013).
- [839] G. Aad (ATLAS), “Measurement of the Azimuthal Angle Dependence of Inclusive Jet Yields in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Rev. Lett. **111**, 152301 (2013).
- [840] G. Aad (ATLAS), “Measurement of the cross-section for W boson production in association with b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **06**, 084 (2013).
- [841] G. Aad (ATLAS), “Measurement of the differential cross-section of B^+ meson production in pp collisions at $\sqrt{s} = 7$ TeV at ATLAS”, JHEP **10**, 042 (2013).
- [842] G. Aad (ATLAS), “Measurement of the distributions of event-by-event flow harmonics in lead-lead collisions at $\sqrt{s} = 2.76$ TeV with the ATLAS detector at the LHC”, JHEP **11**, 183 (2013).
- [843] G. Aad (ATLAS), “Measurement of the flavour composition of dijet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **73**, 2301 (2013).
- [844] G. Aad (ATLAS), “Measurement of the high-mass Drell–Yan differential cross-section in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. B **725**, 223–242 (2013).
- [845] G. Aad (ATLAS), “Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s}=2.76$ TeV and comparison to the inclusive jet cross section at $\sqrt{s}=7$ TeV using the ATLAS detector”, Eur. Phys. J. C **73**, 2509 (2013).
- [846] G. Aad (ATLAS), “Measurement of the jet radius and transverse momentum dependence of inclusive jet suppression in lead-lead collisions at $\sqrt{s_{NN}}=2.76$ TeV with the ATLAS detector”, Phys. Lett. B **719**, 220–241 (2013).
- [847] G. Aad (ATLAS), “Measurement of the production cross section of jets in association with a Z boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **07**, 032 (2013).
- [848] G. Aad (ATLAS), “Measurement of the top quark charge in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **11**, 031 (2013).
- [849] G. Aad (ATLAS), “Measurement of the ttbar production cross section in the tau+jets channel using the ATLAS detector”, Eur. Phys. J. C **73**, 2328 (2013).

- [850] G. Aad (ATLAS), “Measurement of Top Quark Polarization in Top-Antitop Events from Proton-Proton Collisions at $\sqrt{s} = 7$ TeV Using the ATLAS Detector”, Phys. Rev. Lett. **111**, 232002 (2013).
- [851] G. Aad (ATLAS), “Measurement of Upsilon production in 7 TeV pp collisions at ATLAS”, Phys. Rev. D **87**, 052004 (2013).
- [852] G. Aad (ATLAS), “Measurement with the ATLAS detector of multi-particle azimuthal correlations in p+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Lett. B **725**, 60–78 (2013).
- [853] G. Aad (ATLAS), “Measurements of $W\gamma$ and $Z\gamma$ production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector at the LHC”, Phys. Rev. D **87**, [Erratum: Phys.Rev.D 91, 119901 (2015)], 112003 (2013).
- [854] G. Aad (ATLAS), “Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC”, Phys. Lett. B **726**, [Erratum: Phys.Lett.B 734, 406–406 (2014)], 88–119 (2013).
- [855] G. Aad (ATLAS), “Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **73**, 2261 (2013).
- [856] G. Aad (ATLAS), “Multi-channel search for squarks and gluinos in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **73**, 2362 (2013).
- [857] G. Aad (ATLAS), “Observation of Associated Near-Side and Away-Side Long-Range Correlations in $\sqrt{s_{NN}}=5.02$ TeV Proton-Lead Collisions with the ATLAS Detector”, Phys. Rev. Lett. **110**, 182302 (2013).
- [858] G. Aad (ATLAS), “Performance of jet substructure techniques for large- R jets in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **09**, 076 (2013).
- [859] G. Aad (ATLAS), “Search for $t\bar{t}$ resonances in the lepton plus jets final state with ATLAS using 4.7 fb $^{-1}$ of pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. D **88**, 012004 (2013).
- [860] G. Aad (ATLAS), “Search for a heavy narrow resonance decaying to $e\mu$, $e\tau$, or $\mu\tau$ with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, Phys. Lett. B **723**, 15–32 (2013).
- [861] G. Aad (ATLAS), “Search for a light charged Higgs boson in the decay channel $H^+ \rightarrow c\bar{s}$ in $t\bar{t}$ events using pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **73**, 2465 (2013).
- [862] G. Aad (ATLAS), “Search for charged Higgs bosons through the violation of lepton universality in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, JHEP **03**, 076 (2013).
- [863] G. Aad (ATLAS), “Search for charginos nearly mass degenerate with the lightest neutralino based on a disappearing-track signature in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector”, Phys. Rev. D **88**, 112006 (2013).
- [864] G. Aad (ATLAS), “Search for contact interactions and large extra dimensions in dilepton events from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **87**, 015010 (2013).
- [865] G. Aad (ATLAS), “Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector”, JHEP **04**, 075 (2013).
- [866] G. Aad (ATLAS), “Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **110**, 011802 (2013).
- [867] G. Aad (ATLAS), “Search for direct chargino production in anomaly-mediated supersymmetry breaking models based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **01**, 131 (2013).
- [868] G. Aad (ATLAS), “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Phys. Lett. B **718**, 841–859 (2013).
- [869] G. Aad (ATLAS), “Search for direct slepton and gaugino production in final states with two leptons and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. B **718**, 879–901 (2013).
- [870] G. Aad (ATLAS), “Search for direct third-generation squark pair production in final states with missing transverse momentum and two b -jets in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, JHEP **10**, 189 (2013).
- [871] G. Aad (ATLAS), “Search for displaced muonic lepton jets from light Higgs boson decay in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **721**, 32–50 (2013).
- [872] G. Aad (ATLAS), “Search for excited electrons and muons in $\sqrt{s}=8$ TeV proton-proton collisions with the ATLAS detector”, New J. Phys. **15**, 093011 (2013).

- [873] G. Aad (ATLAS), “Search for Extra Dimensions in diphoton events using proton-proton collisions recorded at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC”, *New J. Phys.* **15**, 043007 (2013).
- [874] G. Aad (ATLAS), “Search for light top squark pair production in final states with leptons and b^- jets with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, *Phys. Lett. B* **720**, 13–31 (2013).
- [875] G. Aad (ATLAS), “Search for long-lived stopped R-hadrons decaying out-of-time with pp collisions using the ATLAS detector”, *Phys. Rev. D* **88**, 112003 (2013).
- [876] G. Aad (ATLAS), “Search for long-lived, heavy particles in final states with a muon and multi-track displaced vertex in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **719**, 280–298 (2013).
- [877] G. Aad (ATLAS), “Search for long-lived, multi-charged particles in pp collisions at $\sqrt{s}=7$ TeV using the ATLAS detector”, *Phys. Lett. B* **722**, 305–323 (2013).
- [878] G. Aad (ATLAS), “Search for microscopic black holes in a like-sign dimuon final state using large track multiplicity with the ATLAS detector”, *Phys. Rev. D* **88**, 072001 (2013).
- [879] G. Aad (ATLAS), “Search for new phenomena in events with three charged leptons at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Rev. D* **87**, 052002 (2013).
- [880] G. Aad (ATLAS), “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum at $\sqrt{s}=8$ TeV proton-proton collisions using the ATLAS experiment”, *JHEP* **10**, [Erratum: *JHEP* 01, 109 (2014)], 130 (2013).
- [881] G. Aad (ATLAS), “Search for new phenomena in the WW to $\ell\nu\ell'\nu'$ final state in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **718**, 860–878 (2013).
- [882] G. Aad (ATLAS), “Search for nonpointing photons in the diphoton and E_T^{miss} final state in $\sqrt{s}=7$ TeV proton-proton collisions using the ATLAS detector”, *Phys. Rev. D* **88**, 012001 (2013).
- [883] G. Aad (ATLAS), “Search for pair production of heavy top-like quarks decaying to a high-pT W boson and a b quark in the lepton plus jets final state at $\sqrt{s}=7$ TeV with the ATLAS detector”, *Phys. Lett. B* **718**, 1284–1302 (2013).
- [884] G. Aad (ATLAS), “Search for pair-produced massive coloured scalars in four-jet final states with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **73**, 2263 (2013).
- [885] G. Aad (ATLAS), “Search for resonances decaying into top-quark pairs using fully hadronic decays in pp collisions with ATLAS at $\sqrt{s} = 7$ TeV”, *JHEP* **01**, 116 (2013).
- [886] G. Aad (ATLAS), “Search for resonant diboson production in the $WW/WZ \rightarrow \ell\nu jj$ decay channels with the ATLAS detector at $\sqrt{s} = 7$ TeV”, *Phys. Rev. D* **87**, 112006 (2013).
- [887] G. Aad (ATLAS), “Search for single b^* -quark production with the ATLAS detector at $\sqrt{s} = 7$ TeV”, *Phys. Lett. B* **721**, 171–189 (2013).
- [888] G. Aad (ATLAS), “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collision data”, *Phys. Rev. D* **87**, 012008 (2013).
- [889] G. Aad (ATLAS), “Search for supersymmetry in events with photons, bottom quarks, and missing transverse momentum in proton–proton collisions at a centre-of-mass energy of 7 TeV with the ATLAS detector”, *Phys. Lett. B* **719**, 261–279 (2013).
- [890] G. Aad (ATLAS), “Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *JHEP* **02**, 095 (2013).
- [891] G. Aad (ATLAS), “Search for third generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *JHEP* **06**, 033 (2013).
- [892] G. Aad (ATLAS), “Search for WH production with a light Higgs boson decaying to prompt electron-jets in proton-proton collisions at $\sqrt{s}=7$ TeV with the ATLAS detector”, *New J. Phys.* **15**, 043009 (2013).
- [893] G. Aad (ATLAS), “Searches for heavy long-lived sleptons and R-Hadrons with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, *Phys. Lett. B* **720**, 277–308 (2013).
- [894] G. Aad (ATLAS), “Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC”, *Eur. Phys. J. C* **73**, 2305 (2013).
- [895] G. Aad (ATLAS), “Technical Design Report for the Phase-I Upgrade of the ATLAS TDAQ System”, (2013).
- [896] G. Aad (ATLAS), “Triggers for displaced decays of long-lived neutral particles in the ATLAS detector”, *JINST* **8**, P07015 (2013).
- [897] T. Aaltonen (CDF), “ W boson polarization measurement in the $t\bar{t}$ dilepton channel using the CDF II Detector”, *Phys. Lett. B* **722**, 48–54 (2013).

- [898] T. Aaltonen (CDF), “Measurement of the B_c^- meson lifetime in the decay $B_c^- \rightarrow J/\psi \pi^-$ ”, Phys. Rev. D **87**, 011101 (2013).
- [899] T. Aaltonen (CDF), “Measurement of the mass difference between top and antitop quarks”, Phys. Rev. D **87**, 052013 (2013).
- [900] T. Aaltonen (CDF), “Observation of the Production of a W Boson in Association with a Single Charm Quark”, Phys. Rev. Lett. **110**, 071801 (2013).
- [901] T. Aaltonen (CDF), “Production of $K_S^0, K^{*\pm}(892)$ and $\phi^0(1020)$ in Minimum Bias Events and K_S^0 and Λ^0 in Jets in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{TeV}$ ”, Phys. Rev. D **88**, 092005 (2013).
- [902] T. Aaltonen (CDF), “Search for a Two-Higgs-Boson Doublet Using a Simplified Model in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **110**, 121801 (2013).
- [903] T. Aaltonen (CDF), “Search for the Higgs boson in the all-hadronic final state using the full CDF data set”, JHEP **02**, 004 (2013).
- [904] T. Aaltonen (CDF), “Searches for the Higgs boson decaying to $W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$ with the CDF II detector”, Phys. Rev. D **88**, 052012 (2013).
- [905] J. Anderson, “A fast hardware tracker for the ATLAS trigger system”, Nucl. Instrum. Meth. A **718**, edited by F. Cervelli, G. Chiarelli, F. Forti, M. Grassi, and A. Scribano, 258–259 (2013).
- [906] G. Aad (ATLAS), “A measurement of the ratio of the W and Z cross sections with exactly one associated jet in pp collisions at $\sqrt{s} = 7\text{ TeV}$ with ATLAS”, Phys. Lett. B **708**, 221–240 (2012).
- [907] G. Aad (ATLAS), “A particle consistent with the Higgs Boson observed with the ATLAS Detector at the Large Hadron Collider”, Science **338**, 1576–1582 (2012).
- [908] G. Aad (ATLAS), “A search for $t\bar{t}$ resonances in lepton+jets events with highly boosted top quarks collected in pp collisions at $\sqrt{s} = 7\text{ TeV}$ with the ATLAS detector”, JHEP **09**, 041 (2012).
- [909] G. Aad (ATLAS), “A search for $t\bar{t}$ resonances with the ATLAS detector in 2.05 fb^{-1} of proton-proton collisions at $\sqrt{s} = 7\text{ TeV}$ ”, Eur. Phys. J. C **72**, 2083 (2012).
- [910] G. Aad (ATLAS), “A search for flavour changing neutral currents in top-quark decays in pp collision data collected with the ATLAS detector at $\sqrt{s} = 7\text{ TeV}$ ”, JHEP **09**, 139 (2012).
- [911] G. Aad (ATLAS), “A study of the material in the ATLAS inner detector using secondary hadronic interactions”, JINST **7**, P01013 (2012).
- [912] G. Aad (ATLAS), “ATLAS Measurements of the Properties of Jets for Boosted Particle Searches”, Phys. Rev. D **86**, 072006 (2012).
- [913] G. Aad (ATLAS), “ATLAS search for a heavy gauge boson decaying to a charged lepton and a neutrino in pp collisions at $\sqrt{s} = 7\text{ TeV}$ ”, Eur. Phys. J. C **72**, 2241 (2012).
- [914] G. Aad (ATLAS), “Combined search for the Standard Model Higgs boson in pp collisions at $\sqrt{s} = 7\text{ TeV}$ with the ATLAS detector”, Phys. Rev. D **86**, 032003 (2012).
- [915] G. Aad (ATLAS), “Combined search for the Standard Model Higgs boson using up to 4.9 fb^{-1} of pp collision data at $\sqrt{s} = 7\text{ TeV}$ with the ATLAS detector at the LHC”, Phys. Lett. B **710**, 49–66 (2012).
- [916] G. Aad (ATLAS), “Determination of the strange quark density of the proton from ATLAS measurements of the $W \rightarrow \ell\nu$ and $Z \rightarrow \ell\ell$ cross sections”, Phys. Rev. Lett. **109**, 012001 (2012).
- [917] G. Aad (ATLAS), “Electron performance measurements with the ATLAS detector using the 2010 LHC proton-proton collision data”, Eur. Phys. J. C **72**, 1909 (2012).
- [918] G. Aad (ATLAS), “Evidence for the associated production of a W boson and a top quark in ATLAS at $\sqrt{s} = 7\text{ TeV}$ ”, Phys. Lett. B **716**, 142–159 (2012).
- [919] G. Aad (ATLAS), “Forward-backward correlations and charged-particle azimuthal distributions in pp interactions using the ATLAS detector”, JHEP **07**, 019 (2012).
- [920] G. Aad (ATLAS), “Further search for supersymmetry at $\sqrt{s} = 7\text{ TeV}$ in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector”, Phys. Rev. D **86**, 092002 (2012).
- [921] G. Aad (ATLAS), “Hunt for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in 4.7 fb^{-1} of $\sqrt{s} = 7\text{ TeV}$ proton-proton collisions”, JHEP **07**, 167 (2012).
- [922] G. Aad (ATLAS), “Jet mass and substructure of inclusive jets in $\sqrt{s} = 7\text{ TeV}$ pp collisions with the ATLAS experiment”, JHEP **05**, 128 (2012).
- [923] G. Aad (ATLAS), “Kshort and Λ production in pp interactions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector at the LHC”, Phys. Rev. D **85**, 012001 (2012).
- [924] G. Aad (ATLAS), “Measurement of τ polarization in $W^- \rightarrow \tau\nu$ decays with the ATLAS detector in pp collisions at $\sqrt{s} = 7\text{ TeV}$ ”, Eur. Phys. J. C **72**, 2062 (2012).
- [925] G. Aad (ATLAS), “Measurement of $D^{*+/-}$ meson production in jets from pp collisions at $\sqrt{s} = 7\text{ TeV}$ with the ATLAS detector”, Phys. Rev. D **85**, 052005 (2012).

- [926] G. Aad (ATLAS), “Measurement of $t\bar{t}$ production with a veto on additional jet activity in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. C **72**, 2043 (2012).
- [927] G. Aad (ATLAS), “Measurement of $W\gamma$ and $Z\gamma$ production cross sections in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous triple gauge couplings with the ATLAS detector”, Phys. Lett. B **717**, 49–69 (2012).
- [928] G. Aad (ATLAS), “Measurement of WZ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **72**, 2173 (2012).
- [929] G. Aad (ATLAS), “Measurement of event shapes at large momentum transfer with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **72**, 2211 (2012).
- [930] G. Aad (ATLAS), “Measurement of inclusive jet and dijet production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. D **86**, 014022 (2012).
- [931] G. Aad (ATLAS), “Measurement of inclusive two-particle angular correlations in pp collisions with the ATLAS detector at the LHC”, JHEP **05**, 157 (2012).
- [932] G. Aad (ATLAS), “Measurement of the t -channel single top-quark production cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **717**, 330–350 (2012).
- [933] G. Aad (ATLAS), “Measurement of the $W \rightarrow \tau\nu_\tau$ cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, Phys. Lett. B **706**, 276–294 (2012).
- [934] G. Aad (ATLAS), “Measurement of the WW cross section in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector and limits on anomalous gauge couplings”, Phys. Lett. B **712**, 289–308 (2012).
- [935] G. Aad (ATLAS), “Measurement of the $W^\pm Z$ production cross section and limits on anomalous triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **709**, 341–357 (2012).
- [936] G. Aad (ATLAS), “Measurement of the ZZ production cross section and limits on anomalous neutral triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **108**, 041804 (2012).
- [937] G. Aad (ATLAS), “Measurement of the azimuthal anisotropy for charged particle production in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector”, Phys. Rev. C **86**, 014907 (2012).
- [938] G. Aad (ATLAS), “Measurement of the azimuthal ordering of charged hadrons with the ATLAS detector”, Phys. Rev. D **86**, 052005 (2012).
- [939] G. Aad (ATLAS), “Measurement of the b-hadron production cross section using decays to $D^*\mu^-X$ final states in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. B **864**, 341–381 (2012).
- [940] G. Aad (ATLAS), “Measurement of the centrality dependence of the charged particle pseudorapidity distribution in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. B **710**, 363–382 (2012).
- [941] G. Aad (ATLAS), “Measurement of the charge asymmetry in top quark pair production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. C **72**, 2039 (2012).
- [942] G. Aad (ATLAS), “Measurement of the cross section for the production of a W boson in association with b^- jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **707**, 418–437 (2012).
- [943] G. Aad (ATLAS), “Measurement of the cross section for top-quark pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector using final states with two high-pt leptons”, JHEP **05**, 059 (2012).
- [944] G. Aad (ATLAS), “Measurement of the cross-section for b -jets produced in association with a Z boson at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **706**, 295–313 (2012).
- [945] G. Aad (ATLAS), “Measurement of the inclusive W^\pm and Z/γ cross sections in the electron and muon decay channels in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **85**, 072004 (2012).
- [946] G. Aad (ATLAS), “Measurement of the isolated di-photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **85**, 012003 (2012).
- [947] G. Aad (ATLAS), “Measurement of the polarisation of W bosons produced with large transverse momentum in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, Eur. Phys. J. C **72**, 2001 (2012).
- [948] G. Aad (ATLAS), “Measurement of the production cross section for Z/γ^* in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **85**, 032009 (2012).

- [949] G. Aad (ATLAS), “Measurement of the production cross section of an isolated photon associated with jets in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **85**, 092014 (2012).
- [950] G. Aad (ATLAS), “Measurement of the pseudorapidity and transverse momentum dependence of the elliptic flow of charged particles in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. B **707**, 330–348 (2012).
- [951] G. Aad (ATLAS), “Measurement of the top quark mass with the template method in the $t\bar{t} \rightarrow$ lepton + jets channel using ATLAS data”, Eur. Phys. J. C **72**, 2046 (2012).
- [952] G. Aad (ATLAS), “Measurement of the top quark pair cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV using final states with an electron or a muon and a hadronically decaying τ lepton”, Phys. Lett. B **717**, 89–108 (2012).
- [953] G. Aad (ATLAS), “Measurement of the top quark pair production cross section in pp collisions at $\sqrt{s} = 7$ TeV in dilepton final states with ATLAS”, Phys. Lett. B **707**, 459–477 (2012).
- [954] G. Aad (ATLAS), “Measurement of the top quark pair production cross-section with ATLAS in the single lepton channel”, Phys. Lett. B **711**, 244–263 (2012).
- [955] G. Aad (ATLAS), “Measurement of the Transverse Momentum Distribution of W Bosons in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Phys. Rev. D **85**, 012005 (2012).
- [956] G. Aad (ATLAS), “Measurement of the W boson polarization in top quark decays with the ATLAS detector”, JHEP **06**, 088 (2012).
- [957] G. Aad (ATLAS), “Measurements of the electron and muon inclusive cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **707**, 438–458 (2012).
- [958] G. Aad (ATLAS), “Measurements of the pseudorapidity dependence of the total transverse energy in proton-proton collisions at $\sqrt{s} = 7$ TeV with ATLAS”, JHEP **11**, 033 (2012).
- [959] G. Aad (ATLAS), “Observation of a new χ_b state in radiative transitions to $\Upsilon(1S)$ and $\Upsilon(2S)$ at ATLAS”, Phys. Rev. Lett. **108**, 152001 (2012).
- [960] G. Aad (ATLAS), “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, Phys. Lett. B **716**, 1–29 (2012).
- [961] G. Aad (ATLAS), “Observation of spin correlation in $t\bar{t}$ events from pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. Lett. **108**, 212001 (2012).
- [962] G. Aad (ATLAS), “Performance of Missing Transverse Momentum Reconstruction in Proton-Proton Collisions at 7 TeV with ATLAS”, Eur. Phys. J. C **72**, 1844 (2012).
- [963] G. Aad (ATLAS), “Performance of the ATLAS Trigger System in 2010”, Eur. Phys. J. C **72**, 1849 (2012).
- [964] G. Aad (ATLAS), “Rapidity gap cross sections measured with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **72**, 1926 (2012).
- [965] G. Aad (ATLAS), “Search for a fermiophobic Higgs boson in the diphoton decay channel with the ATLAS detector”, Eur. Phys. J. C **72**, 2157 (2012).
- [966] G. Aad (ATLAS), “Search for a heavy Standard Model Higgs boson in the channel $H \rightarrow ZZ \rightarrow \ell^+ \ell^q \bar{q}$ using the ATLAS detector”, Phys. Lett. B **707**, 27–45 (2012).
- [967] G. Aad (ATLAS), “Search for a heavy top-quark partner in final states with two leptons with the ATLAS detector at the LHC”, JHEP **11**, 094 (2012).
- [968] G. Aad (ATLAS), “Search for a light Higgs boson decaying to long-lived weakly-interacting particles in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **108**, 251801 (2012).
- [969] G. Aad (ATLAS), “Search for a standard model Higgs boson in the mass range 200–600 GeV in the $H \rightarrow ZZ \rightarrow \ell^+ \ell^- q \bar{q}$ decay channel with the ATLAS detector”, Phys. Lett. B **717**, 70–88 (2012).
- [970] G. Aad (ATLAS), “Search for a standard model Higgs boson in the $H \rightarrow ZZ \rightarrow \ell^+ \ell^- \nu \bar{\nu}$ decay channel using 4.7 fb $^{-1}$ of $\sqrt{s} = 7$ TeV data with the ATLAS detector”, Phys. Lett. B **717**, 29–48 (2012).
- [971] G. Aad (ATLAS), “Search for a supersymmetric partner to the top quark in final states with jets and missing transverse momentum at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **109**, 211802 (2012).
- [972] G. Aad (ATLAS), “Search for anomalous production of prompt like-sign lepton pairs at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **12**, 007 (2012).
- [973] G. Aad (ATLAS), “Search for anomalous production of prompt like-sign muon pairs and constraints on physics beyond the Standard Model with the ATLAS detector”, Phys. Rev. D **85**, 032004 (2012).

- [974] G. Aad (ATLAS), “Search for anomaly-mediated supersymmetry breaking with the ATLAS detector based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **72**, 1993 (2012).
- [975] G. Aad (ATLAS), “Search for charged Higgs bosons decaying via $H^+ \rightarrow \tau\nu$ in top quark pair events using pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *JHEP* **06**, 039 (2012).
- [976] G. Aad (ATLAS), “Search for contact interactions in dilepton events from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **712**, 40–58 (2012).
- [977] G. Aad (ATLAS), “Search for decays of stopped, long-lived particles from 7 TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **72**, 1965 (2012).
- [978] G. Aad (ATLAS), “Search for diphoton events with large missing transverse momentum in 7 TeV proton-proton collision data with the ATLAS detector”, *Phys. Lett. B* **718**, 411–430 (2012).
- [979] G. Aad (ATLAS), “Search for diphoton events with large missing transverse momentum in fb^{-1} of 7 TeV proton-proton collision data with the ATLAS detector”, *Phys. Lett. B* **710**, 519–537 (2012).
- [980] G. Aad (ATLAS), “Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions using 4.7 fb^{-1} of ATLAS data”, *Phys. Rev. Lett.* **109**, 211803 (2012).
- [981] G. Aad (ATLAS), “Search for displaced vertices arising from decays of new heavy particles in 7 TeV pp collisions at ATLAS”, *Phys. Lett. B* **707**, 478–496 (2012).
- [982] G. Aad (ATLAS), “Search for doubly-charged Higgs bosons in like-sign dilepton final states at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **72**, 2244 (2012).
- [983] G. Aad (ATLAS), “Search for down-type fourth generation quarks with the ATLAS detector in events with one lepton and hadronically decaying W bosons”, *Phys. Rev. Lett.* **109**, 032001 (2012).
- [984] G. Aad (ATLAS), “Search for events with large missing transverse momentum, jets, and at least two tau leptons in 7 TeV proton-proton collision data with the ATLAS detector”, *Phys. Lett. B* **714**, 180–196 (2012).
- [985] G. Aad (ATLAS), “Search for excited leptons in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Rev. D* **85**, 072003 (2012).
- [986] G. Aad (ATLAS), “Search for extra dimensions using diphoton events in 7 TeV proton-proton collisions with the ATLAS detector”, *Phys. Lett. B* **710**, 538–556 (2012).
- [987] G. Aad (ATLAS), “Search for FCNC single top-quark production at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **712**, 351–369 (2012).
- [988] G. Aad (ATLAS), “Search for first generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **709**, [Erratum: *Phys. Lett. B* 711, 442–455 (2012)], 158–176 (2012).
- [989] G. Aad (ATLAS), “Search for gluinos in events with two same-sign leptons, jets and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, *Phys. Rev. Lett.* **108**, 241802 (2012).
- [990] G. Aad (ATLAS), “Search for heavy neutrinos and right-handed W bosons in events with two leptons and jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **72**, 2056 (2012).
- [991] G. Aad (ATLAS), “Search for heavy vector-like quarks coupling to light quarks in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **712**, 22–39 (2012).
- [992] G. Aad (ATLAS), “Search for high-mass resonances decaying to dilepton final states in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *JHEP* **11**, 138 (2012).
- [993] G. Aad (ATLAS), “Search for lepton flavour violation in the $e\mu$ continuum with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, *Eur. Phys. J. C* **72**, 2040 (2012).
- [994] G. Aad (ATLAS), “Search for light scalar top quark pair production in final states with two leptons with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, *Eur. Phys. J. C* **72**, 2237 (2012).
- [995] G. Aad (ATLAS), “Search for magnetic monopoles in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, *Phys. Rev. Lett.* **109**, 261803 (2012).
- [996] G. Aad (ATLAS), “Search for new particles decaying to ZZ using final states with leptons and jets with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, *Phys. Lett. B* **712**, 331–350 (2012).
- [997] G. Aad (ATLAS), “Search for New Phenomena in $t\bar{t}$ Events With Large Missing Transverse Momentum in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **108**, 041805 (2012).
- [998] G. Aad (ATLAS), “Search for New Physics in the Dijet Mass Distribution using 1 fb^{-1} of pp Collision Data at $\sqrt{s} = 7$ TeV collected by the ATLAS Detector”, *Phys. Lett. B* **708**, 37–54 (2012).

- [999] G. Aad (ATLAS), “Search for pair production of a heavy up-type quark decaying to a W boson and a b quark in the lepton+jets channel with the ATLAS detector”, Phys. Rev. Lett. **108**, 261802 (2012).
- [1000] G. Aad (ATLAS), “Search for pair production of a new quark that decays to a Z boson and a bottom quark with the ATLAS detector”, Phys. Rev. Lett. **109**, 071801 (2012).
- [1001] G. Aad (ATLAS), “Search for pair production of massive particles decaying into three quarks with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, JHEP **12**, 086 (2012).
- [1002] G. Aad (ATLAS), “Search for pair-produced heavy quarks decaying to Wq in the two-lepton channel at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **86**, 012007 (2012).
- [1003] G. Aad (ATLAS), “Search for production of resonant states in the photon-jet mass distribution using pp collisions at $\sqrt{s} = 7$ TeV collected by the ATLAS detector”, Phys. Rev. Lett. **108**, 211802 (2012).
- [1004] G. Aad (ATLAS), “Search for R-parity-violating supersymmetry in events with four or more leptons in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, JHEP **12**, 124 (2012).
- [1005] G. Aad (ATLAS), “Search for resonant WZ production in the $WZ \rightarrow \ell\nu\ell'\ell'$ channel in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Phys. Rev. D **85**, 112012 (2012).
- [1006] G. Aad (ATLAS), “Search for resonant top plus jet production in $t\bar{t} +$ jets events with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. D **86**, 091103 (2012).
- [1007] G. Aad (ATLAS), “Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **04**, 069 (2012).
- [1008] G. Aad (ATLAS), “Search for scalar bottom quark pair production with the ATLAS detector in pp Collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. Lett. **108**, 181802 (2012).
- [1009] G. Aad (ATLAS), “Search for scalar top quark pair production in natural gauge mediated supersymmetry models with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. B **715**, 44–60 (2012).
- [1010] G. Aad (ATLAS), “Search for second generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **72**, 2151 (2012).
- [1011] G. Aad (ATLAS), “Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. B **710**, 67–85 (2012).
- [1012] G. Aad (ATLAS), “Search for strong gravity signatures in same-sign dimuon final states using the ATLAS detector at the LHC”, Phys. Lett. B **709**, 322–340 (2012).
- [1013] G. Aad (ATLAS), “Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b^- jets with the ATLAS detector”, Phys. Rev. D **85**, 112006 (2012).
- [1014] G. Aad (ATLAS), “Search for Supersymmetry in Events with Large Missing Transverse Momentum, Jets, and at Least One Tau Lepton in 7 TeV Proton-Proton Collision Data with the ATLAS Detector”, Eur. Phys. J. C **72**, 2215 (2012).
- [1015] G. Aad (ATLAS), “Search for supersymmetry in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Phys. Rev. Lett. **108**, 261804 (2012).
- [1016] G. Aad (ATLAS), “Search for supersymmetry in final states with jets, missing transverse momentum and one isolated lepton in $\sqrt{s} = 7$ TeV pp collisions using 1 fb^{-1} of ATLAS data”, Phys. Rev. D **85**, [Erratum: Phys.Rev.D 87, 099903 (2013)], 012006 (2012).
- [1017] G. Aad (ATLAS), “Search for supersymmetry with jets, missing transverse momentum and at least one hadronically decaying τ lepton in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **714**, 197–214 (2012).
- [1018] G. Aad (ATLAS), “Search for tb resonances in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **109**, 081801 (2012).
- [1019] G. Aad (ATLAS), “Search for TeV-scale gravity signatures in final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 7$ TeV”, Phys. Lett. B **716**, 122–141 (2012).
- [1020] G. Aad (ATLAS), “Search for the decay $B_s^0 \rightarrow \mu\mu$ with the ATLAS detector”, Phys. Lett. B **713**, 387–407 (2012).
- [1021] G. Aad (ATLAS), “Search for the Higgs boson in the $H \rightarrow WW(*) \rightarrow \ell\nu\ell\nu$ decay channel in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **108**, 111802 (2012).
- [1022] G. Aad (ATLAS), “Search for the Higgs boson in the $H \rightarrow WW \rightarrow l\nu jj$ decay channel at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **718**, 391–410 (2012).
- [1023] G. Aad (ATLAS), “Search for the Standard Model Higgs boson in the $H \rightarrow WW(*) \rightarrow \ell\nu\ell\nu$ decay mode with 4.7 /fb of ATLAS data at $\sqrt{s} = 7$ TeV”, Phys. Lett. B **716**, 62–81 (2012).

- [1024] G. Aad (ATLAS), “Search for the Standard Model Higgs boson in the H to $\tau^+\tau^-$ decay mode in $\sqrt{s} = 7$ TeV pp collisions with ATLAS”, JHEP **09**, 070 (2012).
- [1025] G. Aad (ATLAS), “Search for the Standard Model Higgs boson in the decay channel $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$ with 4.8 fb^{-1} of pp collision data at $\sqrt{s} = 7$ TeV with ATLAS”, Phys. Lett. B **710**, 383–402 (2012).
- [1026] G. Aad (ATLAS), “Search for the Standard Model Higgs boson in the diphoton decay channel with 4.9 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV with ATLAS”, Phys. Rev. Lett. **108**, 111803 (2012).
- [1027] G. Aad (ATLAS), “Search for the Standard Model Higgs boson produced in association with a vector boson and decaying to a b -quark pair with the ATLAS detector”, Phys. Lett. B **718**, 369–390 (2012).
- [1028] G. Aad (ATLAS), “Search for top and bottom squarks from gluino pair production in final states with missing transverse energy and at least three b -jets with the ATLAS detector”, Eur. Phys. J. C **72**, 2174 (2012).
- [1029] G. Aad (ATLAS), “Searches for supersymmetry with the ATLAS detector using final states with two leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. B **709**, 137–157 (2012).
- [1030] G. Aad (ATLAS), “Study of jets produced in association with a W boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **85**, 092002 (2012).
- [1031] G. Aad (ATLAS), “Time-dependent angular analysis of the decay $B_s^0 \rightarrow J/\psi\phi$ and extraction of $\Delta\Gamma_s$ and the CP-violating weak phase ϕ_s by ATLAS”, JHEP **12**, 072 (2012).
- [1032] G. Aad (ATLAS), “Underlying event characteristics and their dependence on jet size of charged-particle jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **86**, 072004 (2012).
- [1033] T. Aaltonen (CDF), “A Search for dark matter in events with one jet and missing transverse energy in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 211804 (2012).
- [1034] T. Aaltonen (CDF), “An additional study of multi-muon events produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Lett. B **710**, 278–283 (2012).
- [1035] T. Aaltonen (CDF, D0), “Combination of CDF and D0 measurements of the W boson helicity in top quark decays”, Phys. Rev. D **85**, 071106 (2012).
- [1036] T. Aaltonen (CDF, D0), “Combination of the top-quark mass measurements from the Tevatron collider”, Phys. Rev. D **86**, 092003 (2012).
- [1037] T. Aaltonen (CDF), “Combined search for the standard model Higgs boson decaying to a bb pair using the full CDF data set”, Phys. Rev. Lett. **109**, 111802 (2012).
- [1038] T. Aaltonen (CDF), “Diffractive Dijet Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **86**, 032009 (2012).
- [1039] T. Aaltonen (CDF, D0), “Evidence for a particle produced in association with weak bosons and decaying to a bottom-antibottom quark pair in Higgs boson searches at the Tevatron”, Phys. Rev. Lett. **109**, 071804 (2012).
- [1040] T. Aaltonen (CDF), “Evidence for the charmless annihilation decay mode $B_s^0 \rightarrow \pi^+\pi^-$ ”, Phys. Rev. Lett. **108**, 211803 (2012).
- [1041] T. Aaltonen (CDF), “Measurement of $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$ Branching Ratios”, Phys. Rev. Lett. **108**, 201801 (2012).
- [1042] T. Aaltonen (CDF), “Measurement of ZZ production in leptonic final states at \sqrt{s} of 1.96 TeV at CDF”, Phys. Rev. Lett. **108**, 101801 (2012).
- [1043] T. Aaltonen (CDF), “Measurement of CP-violating asymmetries in $D^0 \rightarrow \pi^+\pi^-$ and $D^0 \rightarrow K^+K^-$ decays at CDF”, Phys. Rev. D **85**, 012009 (2012).
- [1044] T. Aaltonen (CDF), “Measurement of CP-violation asymmetries in $D^0 \rightarrow K_S\pi^+\pi^-$ ”, Phys. Rev. D **86**, 032007 (2012).
- [1045] T. Aaltonen (CDF), “Measurement of the WZ Cross Section and Triple Gauge Couplings in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **86**, 031104 (2012).
- [1046] T. Aaltonen (CDF), “Measurement of the Bottom-Strange Meson Mixing Phase in the Full CDF Data Set”, Phys. Rev. Lett. **109**, 171802 (2012).
- [1047] T. Aaltonen (CDF), “Measurement of the branching fraction $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-\pi^+\pi^-)$ at CDF”, Phys. Rev. D **85**, 032003 (2012).
- [1048] T. Aaltonen (CDF), “Measurement of the CP-Violating Phase $\beta_s^{J/\psi\phi}$ in $B_s^0 \rightarrow J/\psi\phi$ Decays with the CDF II Detector”, Phys. Rev. D **85**, 072002 (2012).
- [1049] T. Aaltonen (CDF), “Measurement of the difference of CP-violating asymmetries in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays at CDF”, Phys. Rev. Lett. **109**, 111801 (2012).

- [1050] T. Aaltonen (CDF), “Measurement of the masses and widths of the bottom baryons Σ_b^{+-} and Σ_b^{*+-} ”, Phys. Rev. D **85**, 092011 (2012).
- [1051] T. Aaltonen (CDF), “Measurement of the Top Quark Mass in the All-Hadronic Mode at CDF”, Phys. Lett. B **714**, 24–31 (2012).
- [1052] T. Aaltonen (CDF), “Measurements of Angular Distributions of Muons From Υ Meson Decays in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 151802 (2012).
- [1053] T. Aaltonen (CDF), “Measurements of the Angular Distributions in the Decays $B \rightarrow K^{(*)}\mu^+\mu^-$ at CDF”, Phys. Rev. Lett. **108**, 081807 (2012).
- [1054] T. Aaltonen (CDF), “Measurements of the Top-quark Mass and the $t\bar{t}$ Cross Section in the Hadronic $\tau +$ Jets Decay Channel at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **109**, 192001 (2012).
- [1055] T. Aaltonen (CDF), “Novel inclusive search for the Higgs boson in the four-lepton final state at CDF”, Phys. Rev. D **86**, 072012 (2012).
- [1056] T. Aaltonen (CDF), “Observation of Exclusive Gamma Gamma Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 081801 (2012).
- [1057] T. Aaltonen (CDF), “Precise measurement of the W -boson mass with the CDF II detector”, Phys. Rev. Lett. **108**, 151803 (2012).
- [1058] T. Aaltonen (CDF), “Precision Top-Quark Mass Measurements at CDF”, Phys. Rev. Lett. **109**, 152003 (2012).
- [1059] T. Aaltonen (CDF), “Production of Λ , $\bar{\Lambda}^0$, Ξ^\pm and Ω^\pm Hyperons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **86**, 012002 (2012).
- [1060] T. Aaltonen (CDF), “Search for $WZ + ZZ$ production with MET + jets with b enhancement at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **85**, 012002 (2012).
- [1061] T. Aaltonen (CDF), “Search for a dark matter candidate produced in association with a single top quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 201802 (2012).
- [1062] T. Aaltonen (CDF), “Search for a heavy particle decaying to a top quark and a light quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 211805 (2012).
- [1063] T. Aaltonen (CDF), “Search for a Heavy Vector Boson Decaying to Two Gluons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **86**, 112002 (2012).
- [1064] T. Aaltonen (CDF), “Search for a Higgs Boson in the Diphoton Final State in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 011801 (2012).
- [1065] T. Aaltonen (CDF), “Search for a Higgs boson in the diphoton final state using the full CDF data set from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Lett. B **717**, 173–181 (2012).
- [1066] T. Aaltonen (CDF), “Search for a low mass Standard Model Higgs boson in the $\tau - \tau$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 181804 (2012).
- [1067] T. Aaltonen (CDF), “Search for anomalous production of multiple leptons in association with W and Z bosons at CDF”, Phys. Rev. D **85**, 092001 (2012).
- [1068] T. Aaltonen (CDF), “Search for heavy metastable particles decaying to jet pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **85**, 012007 (2012).
- [1069] T. Aaltonen (CDF), “Search for Higgs Bosons Produced in Association with b -quarks”, Phys. Rev. D **85**, 032005 (2012).
- [1070] T. Aaltonen (CDF), “Search for high-mass resonances decaying into ZZ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **85**, 012008 (2012).
- [1071] T. Aaltonen (CDF, D0), “Search for Neutral Higgs Bosons in Events with Multiple Bottom Quarks at the Tevatron”, Phys. Rev. D **86**, 091101 (2012).
- [1072] T. Aaltonen (CDF), “Search for new phenomena in events with two Z bosons and missing transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **85**, 011104 (2012).
- [1073] T. Aaltonen (CDF), “Search for Scalar Top Quark Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, JHEP **10**, 158 (2012).
- [1074] T. Aaltonen (CDF), “Search for Standard Model Higgs Boson Production in Association with a W Boson at CDF”, Phys. Rev. D **85**, 052002 (2012).
- [1075] T. Aaltonen (CDF), “Search for Standard Model Higgs Boson Production in Association with a W Boson Using a Matrix Element Technique at CDF in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **85**, 072001 (2012).
- [1076] T. Aaltonen (CDF), “Search for the Rare Radiative Decay: $W \rightarrow \pi\gamma$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **85**, 032001 (2012).

- [1077] T. Aaltonen (CDF), “Search for the standard model Higgs boson decaying to a $b\bar{b}$ pair in events with no charged leptons and large missing transverse energy using the full CDF data set”, Phys. Rev. Lett. **109**, 111805 (2012).
- [1078] T. Aaltonen (CDF), “Search for the standard model Higgs boson decaying to a $b\bar{b}$ pair in events with one charged lepton and large missing transverse energy using the full CDF data set”, Phys. Rev. Lett. **109**, 111804 (2012).
- [1079] T. Aaltonen (CDF), “Search for the standard model Higgs boson decaying to a $b\bar{b}$ pair in events with two oppositely-charged leptons using the full CDF data set”, Phys. Rev. Lett. **109**, 111803 (2012).
- [1080] T. Aaltonen (CDF), “Search for the standard model Higgs boson produced in association with a W^\pm boson with 7.5 fb^{-1} integrated luminosity at CDF”, Phys. Rev. D **86**, 032011 (2012).
- [1081] T. Aaltonen (CDF), “Search for the standard model Higgs boson produced in association with a Z Boson in 7.9 fb^{-1} of p anti- p collisions at $\sqrt{s} = 1.96 \text{ TeV}$ using the CDF II detector”, Phys. Lett. B **715**, 98–104 (2012).
- [1082] T. Aaltonen (CDF), “Search for the standard model Higgs boson produced in association with top quarks using the full CDF data set”, Phys. Rev. Lett. **109**, 181802 (2012).
- [1083] T. Aaltonen (CDF), “Study of Substructure of High Transverse Momentum Jets Produced in Proton-Antiproton Collisions at $\sqrt{s} = 1.96 \text{ TeV}$ ”, Phys. Rev. D **85**, 091101 (2012).
- [1084] T. Aaltonen (CDF), “Transverse momentum cross section of e^+e^- pairs in the Z -boson region from $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ ”, Phys. Rev. D **86**, 052010 (2012).
- [1085] J. Anderson, “FTK: A Fast Track Trigger for ATLAS”, JINST **7**, edited by F. Palla, C10002 (2012).
- [1086] A. Andreani, “The Fast Tracker real time processor and its impact on muon isolation, tau and b-Jet online selections at ATLAS”, IEEE Trans. Nucl. Sci. **59**, 348–357 (2012).
- [1087] J. D. Hobbs, **M. S. Neubauer**, S. Willenbrock, “Tests of the Standard Electroweak Model at the Energy Frontier”, Rev. Mod. Phys. **84**, 1477–1526 (2012).
- [1088] G. Aad (ATLAS), “Charged-particle multiplicities in pp interactions measured with the ATLAS detector at the LHC”, New J. Phys. **13**, 053033 (2011).
- [1089] G. Aad (ATLAS), “Measurement of Inclusive Jet and Dijet Cross Sections in Proton-Proton Collisions at 7 TeV Centre-of-Mass Energy with the ATLAS Detector”, Eur. Phys. J. C **71**, 1512 (2011).
- [1090] G. Aad (ATLAS), “Measurement of the centrality dependence of J/ψ yields and observation of Z production in lead–lead collisions with the ATLAS detector at the LHC”, Phys. Lett. B **697**, 294–312 (2011).
- [1091] G. Aad (ATLAS), “Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Phys. Rev. D **83**, 052005 (2011).
- [1092] G. Aad (ATLAS), “Search for Quark Contact Interactions in Dijet Angular Distributions in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$ Measured with the ATLAS Detector”, Phys. Lett. B **694**, 327–345 (2011).
- [1093] G. Aad (ATLAS), “Studies of the performance of the ATLAS detector using cosmic-ray muons”, Eur. Phys. J. C **71**, 1593 (2011).
- [1094] G. Aad (ATLAS), “Study of Jet Shapes in Inclusive Jet Production in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$ using the ATLAS Detector”, Phys. Rev. D **83**, 052003 (2011).
- [1095] G. Aad (ATLAS), “Inclusive search for same-sign dilepton signatures in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, JHEP **10**, 107 (2011).
- [1096] G. Aad (ATLAS), “Limits on the production of the Standard Model Higgs Boson in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Eur. Phys. J. C **71**, 1728 (2011).
- [1097] G. Aad (ATLAS), “Luminosity Determination in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$ Using the ATLAS Detector at the LHC”, Eur. Phys. J. C **71**, 1630 (2011).
- [1098] G. Aad (ATLAS), “Measurement of Dijet Azimuthal Decorrelations in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$ ”, Phys. Rev. Lett. **106**, 172002 (2011).
- [1099] G. Aad (ATLAS), “Measurement of dijet production with a veto on additional central jet activity in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ using the ATLAS detector”, JHEP **09**, 053 (2011).
- [1100] G. Aad (ATLAS), “Measurement of multi-jet cross sections in proton-proton collisions at a 7 TeV center-of-mass energy”, Eur. Phys. J. C **71**, 1763 (2011).
- [1101] G. Aad (ATLAS), “Measurement of the $\Upsilon(1S)$ production cross-section in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ in ATLAS”, Phys. Lett. B **705**, 9–27 (2011).
- [1102] G. Aad (ATLAS), “Measurement of the WW cross section in $\sqrt{s} = 7 \text{ TeV}$ pp collisions with ATLAS”, Phys. Rev. Lett. **107**, 041802 (2011).

- [1103] G. Aad (ATLAS), “Measurement of the W charge asymmetry in the $W \rightarrow \mu\nu$ decay mode in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **701**, 31–49 (2011).
- [1104] G. Aad (ATLAS), “Measurement of the differential cross-sections of inclusive, prompt and non-prompt J/ψ production in proton-proton collisions at $\sqrt{s} = 7$ TeV”, Nucl. Phys. B **850**, 387–444 (2011).
- [1105] G. Aad (ATLAS), “Measurement of the inclusive and dijet cross-sections of b^- jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. C **71**, 1846 (2011).
- [1106] G. Aad (ATLAS), “Measurement of the inclusive isolated prompt photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV using 35 pb $^{-1}$ of ATLAS data”, Phys. Lett. B **706**, 150–167 (2011).
- [1107] G. Aad (ATLAS), “Measurement of the Inelastic Proton-Proton Cross-Section at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Nature Commun. **2**, 463 (2011).
- [1108] G. Aad (ATLAS), “Measurement of the jet fragmentation function and transverse profile in proton-proton collisions at a center-of-mass energy of 7 TeV with the ATLAS detector”, Eur. Phys. J. C **71**, 1795 (2011).
- [1109] G. Aad (ATLAS), “Measurement of the production cross section for W-bosons in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **698**, 325–345 (2011).
- [1110] G. Aad (ATLAS), “Measurement of the top quark-pair production cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **71**, 1577 (2011).
- [1111] G. Aad (ATLAS), “Measurement of the transverse momentum distribution of Z/γ^* bosons in proton-proton collisions at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. B **705**, 415–434 (2011).
- [1112] G. Aad (ATLAS), “Measurement of the Z to tau tau Cross Section with the ATLAS Detector”, Phys. Rev. D **84**, 112006 (2011).
- [1113] G. Aad (ATLAS), “Measurement of underlying event characteristics using charged particles in pp collisions at $\sqrt{s} = 900\text{GeV}$ and 7 TeV with the ATLAS detector”, Phys. Rev. D **83**, 112001 (2011).
- [1114] G. Aad (ATLAS), “Measurement of $W\gamma$ and $Z\gamma$ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, JHEP **09**, 072 (2011).
- [1115] G. Aad (ATLAS), “Measurements of underlying-event properties using neutral and charged particles in pp collisions at 900 GeV and 7 TeV with the ATLAS detector at the LHC”, Eur. Phys. J. C **71**, 1636 (2011).
- [1116] G. Aad (ATLAS), “Properties of jets measured from tracks in proton-proton collisions at center-of-mass energy $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. D **84**, 054001 (2011).
- [1117] G. Aad (ATLAS), “Search for a heavy gauge boson decaying to a charged lepton and a neutrino in 1 fb $^{-1}$ of pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Lett. B **705**, 28–46 (2011).
- [1118] G. Aad (ATLAS), “Search for a heavy neutral particle decaying into an electron and a muon using 1 fb $^{-1}$ of ATLAS data”, Eur. Phys. J. C **71**, 1809 (2011).
- [1119] G. Aad (ATLAS), “Search for a heavy particle decaying into an electron and a muon with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, Phys. Rev. Lett. **106**, 251801 (2011).
- [1120] G. Aad (ATLAS), “Search for a Standard Model Higgs boson in the $H \rightarrow ZZ \rightarrow l^+l^- \nu\nu^-$ decay channel with the ATLAS detector”, Phys. Rev. Lett. **107**, 221802 (2011).
- [1121] G. Aad (ATLAS), “Search for an excess of events with an identical flavour lepton pair and significant missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS detector”, Eur. Phys. J. C **71**, 1647 (2011).
- [1122] G. Aad (ATLAS), “Search for Contact Interactions in Dimuon Events from pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Phys. Rev. D **84**, 011101 (2011).
- [1123] G. Aad (ATLAS), “Search for dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **107**, 272002 (2011).
- [1124] G. Aad (ATLAS), “Search for Diphoton Events with Large Missing Transverse Energy in 7 TeV Proton-Proton Collisions with the ATLAS Detector”, Phys. Rev. Lett. **106**, 121803 (2011).
- [1125] G. Aad (ATLAS), “Search for Diphoton Events with Large Missing Transverse Energy with 36 pb $^{-1}$ of 7 TeV Proton-Proton Collision Data with the ATLAS Detector”, Eur. Phys. J. C **71**, 1744 (2011).
- [1126] G. Aad (ATLAS), “Search for heavy long-lived charged particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. B **703**, 428–446 (2011).
- [1127] G. Aad (ATLAS), “Search for high mass dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, Phys. Lett. B **700**, 163–180 (2011).
- [1128] G. Aad (ATLAS), “Search for high-mass states with one lepton plus missing transverse momentum in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. B **701**, 50–69 (2011).

- [1129] G. Aad (ATLAS), “Search for Massive Colored Scalars in Four-Jet Final States in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS Detector”, *Eur. Phys. J. C* **71**, 1828 (2011).
- [1130] G. Aad (ATLAS), “Search for Massive Long-lived Highly Ionising Particles with the ATLAS Detector at the LHC”, *Phys. Lett. B* **698**, 353–370 (2011).
- [1131] G. Aad (ATLAS), “Search for neutral MSSM Higgs bosons decaying to $\tau^+\tau^-$ pairs in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Lett. B* **705**, 174–192 (2011).
- [1132] G. Aad (ATLAS), “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, *JHEP* **11**, 099 (2011).
- [1133] G. Aad (ATLAS), “Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, *Phys. Lett. B* **705**, 294–312 (2011).
- [1134] G. Aad (ATLAS), “Search for New Physics in Dijet Mass and Angular Distributions in pp Collisions at $\sqrt{s} = 7$ TeV Measured with the ATLAS Detector”, *New J. Phys.* **13**, 053044 (2011).
- [1135] G. Aad (ATLAS), “Search for pair production of first or second generation leptoquarks in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector at the LHC”, *Phys. Rev. D* **83**, 112006 (2011).
- [1136] G. Aad (ATLAS), “Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, *Phys. Lett. B* **701**, 186–203 (2011).
- [1137] G. Aad (ATLAS), “Search for stable hadronising squarks and gluinos with the ATLAS experiment at the LHC”, *Phys. Lett. B* **701**, 1–19 (2011).
- [1138] G. Aad (ATLAS), “Search for supersymmetric particles in events with lepton pairs and large missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS experiment”, *Eur. Phys. J. C* **71**, 1682 (2011).
- [1139] G. Aad (ATLAS), “Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b -jets”, *Phys. Lett. B* **701**, 398–416 (2011).
- [1140] G. Aad (ATLAS), “Search for supersymmetry using final states with one lepton, jets, and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV pp ”, *Phys. Rev. Lett.* **106**, 131802 (2011).
- [1141] G. Aad (ATLAS), “Search for the Higgs boson in the $H \rightarrow WW \rightarrow lvjj$ decay channel in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **107**, 231801 (2011).
- [1142] G. Aad (ATLAS), “Search for the Standard Model Higgs boson in the decay channel $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$ with the ATLAS detector”, *Phys. Lett. B* **705**, 435–451 (2011).
- [1143] G. Aad (ATLAS), “Search for the Standard Model Higgs boson in the two photon decay channel with the ATLAS detector at the LHC”, *Phys. Lett. B* **705**, 452–470 (2011).
- [1144] T. Aaltonen (CDF), “A Search for resonant production of $t\bar{t}$ pairs in 4.8 fb^{-1} of integrated luminosity of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. D* **84**, 072004 (2011).
- [1145] T. Aaltonen (CDF), “Evidence for $t\bar{t}\gamma$ Production and Measurement of $\sigma_{t\bar{t}\gamma}/\sigma_{t\bar{t}}$ ”, *Phys. Rev. D* **84**, 031104 (2011).
- [1146] T. Aaltonen (CDF), “Evidence for a Mass Dependent Forward-Backward Asymmetry in Top Quark Pair Production”, *Phys. Rev. D* **83**, 112003 (2011).
- [1147] T. Aaltonen (CDF), “First Measurement of the Angular Coefficients of Drell-Yan e^+e^- pairs in the Z Mass Region from $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. Lett.* **106**, 241801 (2011).
- [1148] T. Aaltonen (CDF), “First Search for Multijet Resonances in $\sqrt{s} = 1.96$ TeV $p\bar{p}$ Collisions”, *Phys. Rev. Lett.* **107**, 042001 (2011).
- [1149] T. Aaltonen (CDF), “Improved determination of the sample composition of dimuon events produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, *Eur. Phys. J. C* **71**, 1720 (2011).
- [1150] T. Aaltonen (CDF), “Invariant Mass Distribution of Jet Pairs Produced in Association with a W boson in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. Lett.* **106**, 171801 (2011).
- [1151] T. Aaltonen (CDF), “Limits on Anomalous Trilinear Gauge Couplings in $Z\gamma$ Events from $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. Lett.* **107**, 051802 (2011).
- [1152] T. Aaltonen (CDF), “Measurement of b Hadron Lifetimes in Exclusive Decays Containing a J/ψ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. Lett.* **106**, 121804 (2011).
- [1153] T. Aaltonen (CDF), “Measurement of $t\bar{t}$ Spin Correlation in $p\bar{p}$ Collisions Using the CDF II Detector at the Tevatron”, *Phys. Rev. D* **83**, 031104 (2011).

- [1154] T. Aaltonen (CDF), “Measurement of branching ratio and B_s^0 lifetime in the decay $B_s^0 \rightarrow J/\psi f_0(980)$ at CDF”, Phys. Rev. D **84**, 052012 (2011).
- [1155] T. Aaltonen (CDF), “Measurement of Event Shapes in Proton-Antiproton Collisions at Center-of-Mass Energy 1.96 TeV”, Phys. Rev. D **83**, 112007 (2011).
- [1156] T. Aaltonen (CDF), “Measurement of Polarization and Search for CP-Violation in $B_s^0 \rightarrow \phi\phi$ Decays”, Phys. Rev. Lett. **107**, 261802 (2011).
- [1157] T. Aaltonen (CDF), “Measurement of the B_s Lifetime in Fully and Partially Reconstructed $B_s \rightarrow D_s^-(\phi\pi^-)X$ Decays in $\bar{p} - p$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 272001 (2011).
- [1158] T. Aaltonen (CDF), “Measurement of the B^- Lifetime using a Simulation Free Approach for Trigger Bias Correction”, Phys. Rev. D **83**, 032008 (2011).
- [1159] T. Aaltonen (CDF), “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using events with large Missing E_T and jets”, Phys. Rev. D **84**, 032003 (2011).
- [1160] T. Aaltonen (CDF), “Measurement of the $t\bar{t}$ Production Cross Section with an *in situ* Calibration of b -jet Identification Efficiency”, Phys. Rev. D **83**, 071102 (2011).
- [1161] T. Aaltonen (CDF), “Measurement of the Cross Section for Prompt Isolated Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **84**, 052006 (2011).
- [1162] T. Aaltonen (CDF), “Measurement of the Cross Section for Prompt Isolated Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 102003 (2011).
- [1163] T. Aaltonen (CDF), “Measurement of the Forward-Backward Asymmetry in the $B \rightarrow K^{(*)}\mu^+\mu^-$ Decay and First Observation of the $B_s^0 \rightarrow \phi\mu^+\mu^-$ Decay”, Phys. Rev. Lett. **106**, 161801 (2011).
- [1164] T. Aaltonen (CDF), “Measurement of the Mass Difference between t and \bar{t} Quarks”, Phys. Rev. Lett. **106**, 152001 (2011).
- [1165] T. Aaltonen (CDF), “Measurement of the Top Pair Production Cross Section in the Lepton + Jets Channel Using a Jet Flavor Discriminant”, Phys. Rev. D **84**, 031101 (2011).
- [1166] T. Aaltonen (CDF), “Measurement of the Top Quark Mass in the Lepton+Jets Channel Using the Lepton Transverse Momentum”, Phys. Lett. B **698**, 371–379 (2011).
- [1167] T. Aaltonen (CDF), “Measurement of the top-quark mass in the lepton+jets channel using a matrix element technique with the CDF II detector”, Phys. Rev. D **84**, 071105 (2011).
- [1168] T. Aaltonen (CDF), “Measurements of branching fraction ratios and CP-asymmetries in suppressed $B^- \rightarrow D(\rightarrow K^+\pi^-)K^-$ and $B^- \rightarrow D(\rightarrow K^+\pi^-)\pi^-$ decays”, Phys. Rev. D **84**, 091504 (2011).
- [1169] T. Aaltonen (CDF), “Measurements of Direct CP Violating Asymmetries in Charmless Decays of Strange Bottom Mesons and Bottom Baryons”, Phys. Rev. Lett. **106**, 181802 (2011).
- [1170] T. Aaltonen (CDF), “Measurements of the properties of $\Lambda_c(2595)$, $\Lambda_c(2625)$, $\Sigma_c(2455)$, and $\Sigma_c(2520)$ baryons”, Phys. Rev. D **84**, 012003 (2011).
- [1171] T. Aaltonen (CDF), “Observation of $B_s^0 \rightarrow J/\psi K^{*0}(892)$ and $B_s^0 \rightarrow J/\psi K_S^0$ Decays”, Phys. Rev. D **83**, 052012 (2011).
- [1172] T. Aaltonen (CDF), “Observation of the Ξ_b^0 Baryon”, Phys. Rev. Lett. **107**, 102001 (2011).
- [1173] T. Aaltonen (CDF), “Observation of the Baryonic Flavor-Changing Neutral Current Decay $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$ ”, Phys. Rev. Lett. **107**, 201802 (2011).
- [1174] T. Aaltonen (CDF), “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B^0 \rightarrow \mu^+\mu^-$ Decays with CDF II”, Phys. Rev. Lett. **107**, [Addendum: Phys.Rev.Lett. 107, 239903 (2011)], 191801 (2011).
- [1175] T. Aaltonen (CDF), “Search for a Heavy Top-Like Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 261801 (2011).
- [1176] T. Aaltonen (CDF), “Search for a New Heavy Gauge Boson W' with Electron + Missing ET Event Signature in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **83**, 031102 (2011).
- [1177] T. Aaltonen (CDF), “Search for a Very Light CP-Odd Higgs Boson in Top Quark Decays from $p\bar{p}$ Collisions at 1.96 TeV”, Phys. Rev. Lett. **107**, 031801 (2011).
- [1178] T. Aaltonen (CDF), “Search for Heavy Bottom-Like Quarks Decaying to an Electron or Muon and Jets in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 141803 (2011).
- [1179] T. Aaltonen (CDF), “Search for High Mass Resonances Decaying to Muon Pairs in $\sqrt{s} = 1.96$ TeV $p\bar{p}$ Collisions”, Phys. Rev. Lett. **106**, 121801 (2011).
- [1180] T. Aaltonen (CDF), “Search for New T' Particles in Final States with Large Jet Multiplicities and Missing Transverse Energy in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 191803 (2011).
- [1181] T. Aaltonen (CDF), “Search for New Dielectron Resonances and Randall-Sundrum Gravitons at the Collider Detector at Fermilab”, Phys. Rev. Lett. **107**, 051801 (2011).

- [1182] T. Aaltonen (CDF), “Search for New Heavy Particles Decaying to $ZZ \rightarrow \ell\ell\ell\ell, \ell\ell jj$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **83**, 112008 (2011).
- [1183] T. Aaltonen (CDF), “Search for new physics in high p_T like-sign dilepton events at CDF-II”, Phys. Rev. Lett. **107**, 181801 (2011).
- [1184] T. Aaltonen (CDF), “Search for Production of Heavy Particles Decaying to Top Quarks and Invisible Particles in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 191801 (2011).
- [1185] T. Aaltonen (CDF), “Search for Randall-Sundrum Gravitons in the Diphoton Channel at CDF”, Phys. Rev. D **83**, 011102 (2011).
- [1186] T. Aaltonen (CDF), “Search for resonant production of $t\bar{t}$ decaying to jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **84**, 072003 (2011).
- [1187] T. Aaltonen (CDF), “Search for the Higgs Boson in the All-Hadronic Final State using the CDF II Detector”, Phys. Rev. D **84**, 052010 (2011).
- [1188] T. Aaltonen (CDF), “Top quark mass measurement using the template method at CDF”, Phys. Rev. D **83**, 111101 (2011).
- [1189] T. Aaltonen (CDF), “Top-quark mass measurement using events with missing transverse energy and jets at CDF”, Phys. Rev. Lett. **107**, 232002 (2011).
- [1190] S. Amerio, “ATLAS FTK: Fast track trigger”, PoS **VERTEX2011**, 040 (2011).
- [1191] **M. S. Neubauer**, “Diboson production at colliders”, Ann. Rev. Nucl. Part. Sci. **61**, 223–250 (2011).
- [1192] **M. S. Neubauer** (ATLAS), “Search for the Standard Model Higgs Boson in the Lepton + Missing Transverse Energy + Jets Final State in ATLAS”, (2011).
- [1193] G. Aad (ATLAS), “Charged-particle multiplicities in pp interactions at $\sqrt{s} = 900$ GeV measured with the ATLAS detector at the LHC”, Phys. Lett. B **688**, 21–42 (2010).
- [1194] G. Aad (ATLAS), “Commissioning of the ATLAS Muon Spectrometer with Cosmic Rays”, Eur. Phys. J. C **70**, 875–916 (2010).
- [1195] G. Aad (ATLAS), “Drift Time Measurement in the ATLAS Liquid Argon Electromagnetic Calorimeter using Cosmic Muons”, Eur. Phys. J. C **70**, 755–785 (2010).
- [1196] G. Aad (ATLAS), “Measurement of the $W \rightarrow \ell\nu$ and $Z/\gamma^* \rightarrow \ell\ell$ Production Cross Sections in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, JHEP **12**, 060 (2010).
- [1197] G. Aad (ATLAS), “Performance of the ATLAS Detector using First Collision Data”, JHEP **09**, 056 (2010).
- [1198] G. Aad (ATLAS), “Readiness of the ATLAS Liquid Argon Calorimeter for LHC Collisions”, Eur. Phys. J. C **70**, 723–753 (2010).
- [1199] G. Aad (ATLAS), “Readiness of the ATLAS Tile Calorimeter for LHC collisions”, Eur. Phys. J. C **70**, 1193–1236 (2010).
- [1200] G. Aad (ATLAS), “Search for New Particles in Two-Jet Final States in 7 TeV Proton-Proton Collisions with the ATLAS Detector at the LHC”, Phys. Rev. Lett. **105**, 161801 (2010).
- [1201] G. Aad (ATLAS), “The ATLAS Inner Detector commissioning and calibration”, Eur. Phys. J. C **70**, 787–821 (2010).
- [1202] G. Aad (ATLAS), “The ATLAS Simulation Infrastructure”, Eur. Phys. J. C **70**, 823–874 (2010).
- [1203] G. Aad (ATLAS), “Observation of a Centrality-Dependent Dijet Asymmetry in Lead-Lead Collisions at $\sqrt{s_{NN}} = 2.77$ TeV with the ATLAS Detector at the LHC”, Phys. Rev. Lett. **105**, 252303 (2010).
- [1204] T. Aaltonen (CDF), “A Search for the Higgs Boson Using Neural Networks in Events with Missing Energy and b-quark Jets in p anti-p Collisions at $s^{*(1/2)} = 1.96$ -TeV”, Phys. Rev. Lett. **104**, 141801 (2010).
- [1205] T. Aaltonen (CDF), “A Study of the associated production of photons and b-quark jets in p p-bar collisions at $s^{*(1/2)} = 1.96$ -TeV”, Phys. Rev. D **81**, 052006 (2010).
- [1206] T. Aaltonen (CDF, D0), “Combination of Tevatron Searches for the Standard Model Higgs Boson in the W^+W^- Decay Mode”, Phys. Rev. Lett. **104**, 061802 (2010).
- [1207] T. Aaltonen (CDF, D0), “Combined Tevatron Upper Limit on $gg \rightarrow H \rightarrow W^+W^-$ and Constraints on the Higgs Boson Mass in Fourth-Generation Fermion Models”, Phys. Rev. D **82**, 011102 (2010).
- [1208] T. Aaltonen (CDF), “Diffractive W and Z Production at the Fermilab Tevatron”, Phys. Rev. D **82**, 112004 (2010).
- [1209] T. Aaltonen (CDF), “Direct Top-Quark Width Measurement CDF”, Phys. Rev. Lett. **105**, 232003 (2010).
- [1210] T. Aaltonen (CDF), “Exclusion of an Exotic Top Quark with $-4/3$ Electric Charge Using Soft Lepton Tagging”, Phys. Rev. Lett. **105**, 101801 (2010).

- [1211] T. Aaltonen (CDF), “First Measurement of the b-jet Cross Section in Events with a W Boson in p anti-p Collisions at $s^{*}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **104**, 131801 (2010).
- [1212] T. Aaltonen (CDF), “First Measurement of the Ratio $\sigma(t\bar{t})/\sigma(Z/\gamma \rightarrow \ell\ell)$ and Precise Extraction of the $t\bar{t}$ Cross Section”, Phys. Rev. Lett. **105**, 012001 (2010).
- [1213] T. Aaltonen (CDF), “Improved Search for a Higgs Boson Produced in Association with $Z \rightarrow \ell^+\ell^-$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 251802 (2010).
- [1214] T. Aaltonen (CDF), “Inclusive Search for Standard Model Higgs Boson Production in the WW Decay Channel using the CDF II Detector”, Phys. Rev. Lett. **104**, 061803 (2010).
- [1215] T. Aaltonen (CDF), “Measurement of the Λ_b Lifetime in $\Lambda_b \rightarrow \Lambda_c^+\pi^-$ Decays in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 102002 (2010).
- [1216] T. Aaltonen (CDF), “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s}=1.96\text{ TeV}$ using Soft Electron b-Tagging”, Phys. Rev. D **81**, 092002 (2010).
- [1217] T. Aaltonen (CDF), “Measurement of the W^+W^- Production Cross Section and Search for Anomalous $WW\gamma$ and WWZ Couplings in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 201801 (2010).
- [1218] T. Aaltonen (CDF), “Measurement of the $WW + WZ$ Production Cross Section Using a Matrix Element Technique in Lepton + Jets Events”, Phys. Rev. D **82**, 112001 (2010).
- [1219] T. Aaltonen (CDF), “Measurement of the Top Pair Production Cross Section in the Dilepton Decay Channel in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. D **82**, 052002 (2010).
- [1220] T. Aaltonen (CDF), “Measurement of the Top Quark Mass and $p\bar{p} \rightarrow t\bar{t}$ Cross Section in the All-Hadronic Mode with the CDFII Detector”, Phys. Rev. D **81**, 052011 (2010).
- [1221] T. Aaltonen (CDF), “Measurement of the Top Quark Mass in the Dilepton Channel Using m_{T2} at CDF”, Phys. Rev. D **81**, 031102 (2010).
- [1222] T. Aaltonen (CDF), “Measurement of the $WW+WZ$ Production Cross Section Using the Lepton+Jets Final State at CDF II”, Phys. Rev. Lett. **104**, 101801 (2010).
- [1223] T. Aaltonen (CDF), “Measurement of W-Boson Polarization in Top-quark Decay in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 042002 (2010).
- [1224] T. Aaltonen (CDF), “Measurement of Z gamma Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. D **82**, 031103 (2010).
- [1225] T. Aaltonen (CDF), “Measurements of Branching Fraction Ratios and CP Asymmetries in $B^\pm \rightarrow D_{CP}K^\pm$ Decays in Hadron Collisions”, Phys. Rev. D **81**, 031105 (2010).
- [1226] T. Aaltonen (CDF), “Measurements of the top-quark mass using charged particle tracking”, Phys. Rev. D **81**, 032002 (2010).
- [1227] T. Aaltonen (CDF), “Observation of Single Top Quark Production and Measurement of $|V_{tb}|$ with CDF”, Phys. Rev. D **82**, 112005 (2010).
- [1228] T. Aaltonen (CDF), “Search for WW and WZ Resonances Decaying to Electron, Missing E_T , and Two Jets in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 241801 (2010).
- [1229] T. Aaltonen (CDF), “Search for Anomalous Production of Events with Two Photons and Additional Energetic Objects at CDF”, Phys. Rev. D **82**, 052005 (2010).
- [1230] T. Aaltonen (CDF), “Search for New Bottomlike Quark Pair Decays $Q\bar{Q} \rightarrow (tW)(\bar{t}W)$ in Same-Charge Dilepton Events”, Phys. Rev. Lett. **104**, 091801 (2010).
- [1231] T. Aaltonen (CDF), “Search for New Color-Octet Vector Particle Decaying to t anti-t in p anti-p Collisions at $s^{*}(1/2) = 1.96\text{-TeV}$ ”, Phys. Lett. B **691**, 183–190 (2010).
- [1232] T. Aaltonen (CDF), “Search for New Physics with a Dijet plus Missing Transverse Energy Signature in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 131801 (2010).
- [1233] T. Aaltonen (CDF), “Search for Pair Production of Supersymmetric Top Quarks in Dilepton Events from p anti-p Collisions at $S^{*}(1/2) = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 251801 (2010).
- [1234] T. Aaltonen (CDF), “Search for R-parity Violating Decays of τ Sneutrinos to $e\mu$, $\mu\tau$, and $e\tau$ Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 191801 (2010).
- [1235] T. Aaltonen (CDF), “Search for Single Top Quark Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ in the Missing Transverse Energy Plus Jets Topology”, Phys. Rev. D **81**, 072003 (2010).
- [1236] T. Aaltonen (CDF), “Search for Supersymmetry with Gauge-Mediated Breaking in Diphoton Events with Missing Transverse Energy at CDF II”, Phys. Rev. Lett. **104**, 011801 (2010).
- [1237] T. Aaltonen (CDF), “Search for Technicolor Particles Produced in Association with a W Boson at CDF”, Phys. Rev. Lett. **104**, 111802 (2010).
- [1238] T. Aaltonen (CDF), “Search for the Production of Scalar Bottom Quarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 081802 (2010).

- [1239] T. Aaltonen (CDF), “Search for the Supersymmetric Partner of the Top Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **82**, 092001 (2010).
- [1240] T. Aaltonen (CDF), “Studying the Underlying Event in Drell-Yan and High Transverse Momentum Jet Production at the Tevatron”, Phys. Rev. D **82**, 034001 (2010).
- [1241] T. Aaltonen (CDF), “Top Quark Mass Measurement in the Lepton + Jets Channel Using a Matrix Element Method and in situ Jet Energy Calibration”, Phys. Rev. Lett. **105**, 252001 (2010).
- [1242] T. Aaltonen (CDF), “Updated Search for the Flavor-Changing Neutral-Current Decay $D^0 \rightarrow \mu^+ \mu^-$ ”, Phys. Rev. D **82**, 091105 (2010).
- [1243] T. A. Aaltonen (CDF), “Measurement of $d\sigma/dy$ of Drell-Yan e^+e^- pairs in the Z Mass Region from $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Lett. B **692**, 232–239 (2010).
- [1244] M. Abolins (ATLAS), “The ATLAS Trigger/DAQ Authorlist, version 4.0”, (2010).
- [1245] A. Andreani, “Enhancement of the ATLAS Trigger System with a Hardware Tracker Finder FTK”, JINST **5**, C12037 (2010).
- [1246] T. Ciodaro (ATLAS TDAQ), “Muon Detection Based on a Hadronic Calorimeter”, edited by R. Keyser, 772–776 (2010).
- [1247] G. Aad (ATLAS), “Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics”, (2009).
- [1248] G. Aad (ATLAS), “Triggering on Long-Lived Neutral Particles in the ATLAS Detector”, (2009).
- [1249] T. Aaltonen (CDF), “A Measurement of the t anti- t Cross Section in p anti- p Collisions at $s^{*(1/2)} = 1.96$ -TeV using Dilepton Events with a Lepton plus Track Selection”, Phys. Rev. D **79**, 112007 (2009).
- [1250] T. Aaltonen (CDF), “A Search for high-mass resonances decaying to dimuons at CDF”, Phys. Rev. Lett. **102**, 091805 (2009).
- [1251] T. Aaltonen (CDF), “A Search for the Associated Production of the Standard-Model Higgs Boson in the All-Hadronic Channel”, Phys. Rev. Lett. **103**, 221801 (2009).
- [1252] T. Aaltonen (CDF), “A Search for the Higgs Boson Produced in Association with $Z \rightarrow \ell^+ \ell^-$ Using the Matrix Element Method at CDF II”, Phys. Rev. D **80**, 071101 (2009).
- [1253] T. Aaltonen (CDF), “Direct Measurement of the W Production Charge Asymmetry in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **102**, 181801 (2009).
- [1254] T. Aaltonen (CDF), “Evidence for a Narrow Near-Threshold Structure in the $J/\psi\phi$ Mass Spectrum in $B^+ \rightarrow J/\psi\phi K^+$ Decays”, Phys. Rev. Lett. **102**, 242002 (2009).
- [1255] T. Aaltonen (CDF), “First Direct Bound on the Total Width of the Top Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **102**, 042001 (2009).
- [1256] T. Aaltonen (CDF), “First Measurement of the Ratio of Branching Fractions $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- \bar{\nu}_\mu) / B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)$ ”, Phys. Rev. D **79**, 032001 (2009).
- [1257] T. Aaltonen (CDF), “First Measurement of the t anti- t Differential Cross Section $d\sigma/dM(t \text{ anti-}t)$ in p anti- p Collisions at $s^{*(1/2)} = 1.96$ -TeV”, Phys. Rev. Lett. **102**, 222003 (2009).
- [1258] T. Aaltonen (CDF), “First observation of $\bar{B}_s^0 \rightarrow D_s^\pm K^\mp$ and measurement of the ratio of branching fractions $B(\bar{B}_s^0 \rightarrow D_s^\pm K^\mp) / B(\bar{B}_s^0 \rightarrow D_s^\pm \pi^-)$ ”, Phys. Rev. Lett. **103**, 191802 (2009).
- [1259] T. Aaltonen (CDF), “First Observation of Electroweak Single Top Quark Production”, Phys. Rev. Lett. **103**, 092002 (2009).
- [1260] T. Aaltonen (CDF), “First Observation of Vector Boson Pairs in a Hadronic Final State at the Tevatron Collider”, Phys. Rev. Lett. **103**, 091803 (2009).
- [1261] T. Aaltonen (CDF), “First simultaneous measurement of the top quark mass in the lepton + jets and dilepton channels at CDF”, Phys. Rev. D **79**, 092005 (2009).
- [1262] T. Aaltonen (CDF), “Global Search for New Physics with 2.0 fb^{-1} at CDF”, Phys. Rev. D **79**, 011101 (2009).
- [1263] T. Aaltonen (CDF), “Inclusive Search for Squark and Gluino Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **102**, 121801 (2009).
- [1264] T. Aaltonen (CDF), “Measurement of Cross Sections for b Jet Production in Events with a Z Boson in p^- anti- p Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **79**, 052008 (2009).
- [1265] T. Aaltonen (CDF), “Measurement of Particle Production and Inclusive Differential Cross Sections in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **79**, [Erratum: Phys.Rev.D 82, 119903 (2010)], 112005 (2009).
- [1266] T. Aaltonen (CDF), “Measurement of Resonance Parameters of Orbitally Excited Narrow B^0 Mesons”, Phys. Rev. Lett. **102**, 102003 (2009).

- [1267] T. Aaltonen (CDF), “Measurement of the k_T Distribution of Particles in Jets Produced in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 232002 (2009).
- [1268] T. Aaltonen (CDF), “Measurement of the $t\bar{t}$ Production Cross Section in 2 fb^{-1} of $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ Using Lepton Plus Jets Events with Soft Muon b-Tagging”, Phys. Rev. D **79**, 052007 (2009).
- [1269] T. Aaltonen (CDF), “Measurement of the b-Hadron Production Cross Section Using Decays to $MU\text{-}d0$ X Final States in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. D **79**, 092003 (2009).
- [1270] T. Aaltonen (CDF), “Measurement of the fraction of $t\bar{t}$ production via gluon-gluon fusion in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **79**, 031101 (2009).
- [1271] T. Aaltonen (CDF), “Measurement of the Inclusive Isolated Prompt Photon Cross Section in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ using the CDF Detector”, Phys. Rev. D **80**, 111106 (2009).
- [1272] T. Aaltonen (CDF), “Measurement of the top quark mass at CDF using the ‘neutrino ϕ weighting’ template method on a lepton plus isolated track sample”, Phys. Rev. D **79**, 072005 (2009).
- [1273] T. Aaltonen (CDF), “Measurement of the Top Quark Mass Using the Invariant Mass of Lepton Pairs in Soft Muon b-tagged Events”, Phys. Rev. D **80**, 051104 (2009).
- [1274] T. Aaltonen (CDF), “Measurement of the top quark mass with dilepton events selected using neuroevolution at CDF”, Phys. Rev. Lett. **102**, 152001 (2009).
- [1275] T. Aaltonen (CDF), “Measurement of W-Boson Helicity Fractions in Top-Quark Decays Using $\cos\theta^*$ ”, Phys. Lett. B **674**, 160–167 (2009).
- [1276] T. Aaltonen (CDF), “Observation of exclusive charmonium production and $\gamma + \gamma$ to $\mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **102**, 242001 (2009).
- [1277] T. Aaltonen (CDF), “Observation of New Charmless Decays of Bottom Hadrons”, Phys. Rev. Lett. **103**, 031801 (2009).
- [1278] T. Aaltonen (CDF), “Observation of the Ω_b^- and Measurement of the Properties of the Ξ_b^- and Ω_b^- ”, Phys. Rev. D **80**, 072003 (2009).
- [1279] T. Aaltonen (CDF), “Precision Measurement of the $X(3872)$ Mass in $J/\psi\pi^+\pi^-$ Decays”, Phys. Rev. Lett. **103**, 152001 (2009).
- [1280] T. Aaltonen (CDF), “Production of $\psi(2S)$ Mesons in p anti-p Collisions at 1.96-TeV ”, Phys. Rev. D **80**, 031103 (2009).
- [1281] T. Aaltonen (CDF), “Search for a Fermiophobic Higgs Boson Decaying into Diphotons in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **103**, 061803 (2009).
- [1282] T. Aaltonen (CDF), “Search for a Higgs Boson Decaying to Two W Bosons at CDF”, Phys. Rev. Lett. **102**, 021802 (2009).
- [1283] T. Aaltonen (CDF), “Search for a Higgs Boson in $WH \rightarrow \ell\nu b\bar{b}$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **103**, 101802 (2009).
- [1284] T. Aaltonen (CDF), “Search for Anomalous Production of Events with a Photon, Jet, b-quark Jet, and Missing Transverse Energy”, Phys. Rev. D **80**, 052003 (2009).
- [1285] T. Aaltonen (CDF), “Search for charged Higgs bosons in decays of top quarks in p anti-p collisions at $s^{**}(1/2) = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **103**, 101803 (2009).
- [1286] T. Aaltonen (CDF), “Search for exclusive Z boson production and observation of high mass $p\bar{p} \rightarrow \gamma\gamma \rightarrow p + \ell\ell + \bar{p}$ events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **102**, 222002 (2009).
- [1287] T. Aaltonen (CDF), “Search for Gluino-Mediated Sbottom Production in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 221801 (2009).
- [1288] T. Aaltonen (CDF), “Search for Hadronic Decays of W and Z Bosons in Photon Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-GeV}$ ”, Phys. Rev. D **80**, 052011 (2009).
- [1289] T. Aaltonen (CDF), “Search for Higgs bosons predicted in two-Higgs-doublet models via decays to tau lepton pairs in 1.96-TeV p anti-p collisions”, Phys. Rev. Lett. **103**, 201801 (2009).
- [1290] T. Aaltonen (CDF), “Search for High-Mass $e^+ e^-$ Resonances in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 031801 (2009).
- [1291] T. Aaltonen (CDF), “Search for Long-Lived Massive Charged Particles in $1.96\text{ TeV } p\bar{p}$ Collisions”, Phys. Rev. Lett. **103**, 021802 (2009).
- [1292] T. Aaltonen (CDF), “Search for Maximal Flavor Violating Scalars in Same-Charge Lepton Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 041801 (2009).
- [1293] T. Aaltonen (CDF), “Search for narrow resonances lighter than Υ mesons”, Eur. Phys. J. C **62**, 319–326 (2009).

- [1294] T. Aaltonen (CDF), “Search for new particles decaying into dijets in proton-antiproton collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **79**, 112002 (2009).
- [1295] T. Aaltonen (CDF), “Search for new physics in the $\mu\mu + e/\mu + \cancel{E}_T$ channel with a low- p_T lepton threshold at the Collider Detector at Fermilab”, Phys. Rev. D **79**, 052004 (2009).
- [1296] T. Aaltonen (CDF), “Search for Standard Model Higgs Boson Production in Association with a W Boson using a Neural Network Discriminant at CDF”, Phys. Rev. D **80**, 012002 (2009).
- [1297] T. Aaltonen (CDF), “Search for the Decays $B_s^0 \rightarrow e^+\mu^-$ and $B_s^0 \rightarrow e^+e^-$ in CDF Run II”, Phys. Rev. Lett. **102**, 201801 (2009).
- [1298] T. Aaltonen (CDF), “Search for the Neutral Current Top Quark Decay $t \rightarrow Zc$ Using Ratio of Z-Boson + 4 Jets to W-Boson + 4 Jets Production”, Phys. Rev. D **80**, 052001 (2009).
- [1299] T. Aaltonen (CDF), “Search for the Production of Narrow t anti-b Resonances in 1.9 fb⁻¹ of p anti-p Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **103**, 041801 (2009).
- [1300] T. Aaltonen (CDF), “Search for the Rare Decays $B^+ \rightarrow \mu^+\mu^-K^+$, $B^0 \rightarrow \mu^+\mu^-K^{*0}$ (892), and $B_s^0 \rightarrow \mu^+\mu^-\phi$ at CDF”, Phys. Rev. D **79**, 011104 (2009).
- [1301] T. Aaltonen (CDF), “Search for top-quark production via flavor-changing neutral currents in W+1 jet events at CDF”, Phys. Rev. Lett. **102**, 151801 (2009).
- [1302] T. Aaltonen (CDF), “Search for WW and WZ Production in Lepton Plus Jets Final State at CDF”, Phys. Rev. D **79**, 112011 (2009).
- [1303] T. Aaltonen (CDF), “Searching the Inclusive Lepton + Photon + Missing E(T) + b-quark Signature for Radiative Top Quark Decay and Non-Standard-Model Processes”, Phys. Rev. D **80**, 011102 (2009).
- [1304] T. Aaltonen (CDF), “Top Quark Mass Measurement in the $t\bar{t}$ All Hadronic Channel using a Matrix Element Technique in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **79**, 072010 (2009).
- [1305] T. Aaltonen (CDF), “Top Quark Mass Measurement in the Lepton plus Jets Channel Using a Modified Matrix Element Method”, Phys. Rev. D **79**, 072001 (2009).
- [1306] M. Abolins (ATLAS), “The ATLAS Trigger/DAQ Authorlist, version 3.0”, (2009).
- [1307] M. Abolins (ATLAS), “The ATLAS Trigger/DAQ Authorlist, version 3.1”, (2009).
- [1308] A. Abulencia (CDF), “Measurement of the Ratio of Branching Fractions $B(B^+ \rightarrow J/\psi\pi^+)/B(B^+ \rightarrow J/\psi K^+)$ ”, Phys. Rev. D **79**, 112003 (2009).
- [1309] A. Annovi, “Development of FTK architecture: a fast hardware track trigger for the ATLAS detector”, (2009).
- [1310] A. Annovi, “The fast tracker architecture for the LHC baseline luminosity”, PoS **EPS-HEP2009**, 136 (2009).
- [1311] **M. Neubauer**, “A Fast Hardware Tracker for the ATLAS Trigger System”, (2009).
- [1312] T. Aaltonen (CDF), “A Direct measurement of the W boson width in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 071801 (2008).
- [1313] T. Aaltonen (CDF), “Cross-section constrained top quark mass measurement from dilepton events at the Tevatron”, Phys. Rev. Lett. **100**, 062005 (2008).
- [1314] T. Aaltonen (CDF), “Evidence for $D^0 - \bar{D}^0$ mixing using the CDF II Detector”, Phys. Rev. Lett. **100**, 121802 (2008).
- [1315] T. Aaltonen (CDF), “First Flavor-Tagged Determination of Bounds on Mixing-Induced CP Violation in $B_s^0 \rightarrow J/\psi\phi$ Decays”, Phys. Rev. Lett. **100**, 161802 (2008).
- [1316] T. Aaltonen (CDF), “First Measurement of ZZ Production in panti-p Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 201801 (2008).
- [1317] T. Aaltonen (CDF), “First measurement of the fraction of top quark pair production through gluon-gluon fusion”, Phys. Rev. D **78**, 111101 (2008).
- [1318] T. Aaltonen (CDF), “First measurement of the production of a W boson in association with a single charm quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 091803 (2008).
- [1319] T. Aaltonen (CDF), “First Observation of the Decay $B_s^0 \rightarrow D_s^- D_s^+$ and Measurement of Its Branching Ratio”, Phys. Rev. Lett. **100**, 021803 (2008).
- [1320] T. Aaltonen (CDF), “First Run II Measurement of the W Boson Mass”, Phys. Rev. D **77**, 112001 (2008).
- [1321] T. Aaltonen (CDF), “Forward-Backward Asymmetry in Top Quark Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **101**, 202001 (2008).
- [1322] T. Aaltonen (CDF), “Limits on the production of narrow $t\bar{t}$ resonances in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **77**, 051102 (2008).

- [1323] T. Aaltonen (CDF), “Measurement of b -jet Shapes in Inclusive Jet Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **78**, 072005 (2008).
- [1324] T. Aaltonen (CDF), “Measurement of correlated $b\bar{b}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1960\text{ GeV}$ ”, Phys. Rev. D **77**, 072004 (2008).
- [1325] T. Aaltonen (CDF), “Measurement of inclusive jet cross-sections in $Z/\gamma^* \rightarrow e^+e^- + \text{jets}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 102001 (2008).
- [1326] T. Aaltonen (CDF), “Measurement of lifetime and decay-width difference in $B_s^0 \rightarrow J/\psi\phi$ decays”, Phys. Rev. Lett. **100**, 121803 (2008).
- [1327] T. Aaltonen (CDF), “Measurement of Ratios of Fragmentation Fractions for Bottom Hadrons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **77**, 072003 (2008).
- [1328] T. Aaltonen (CDF), “Measurement of the cross section for W^- boson production in association with jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **77**, 011108 (2008).
- [1329] T. Aaltonen (CDF), “Measurement of the Inclusive Jet Cross Section at the Fermilab Tevatron p anti- p Collider Using a Cone-Based Jet Algorithm”, Phys. Rev. D **78**, [Erratum: Phys.Rev.D 79, 119902 (2009)], 052006 (2008).
- [1330] T. Aaltonen (CDF), “Measurement of the Single Top Quark Production Cross Section at CDF”, Phys. Rev. Lett. **101**, 252001 (2008).
- [1331] T. Aaltonen (CDF), “Model-Independent and Quasi-Model-Independent Search for New Physics at CDF”, Phys. Rev. D **78**, 012002 (2008).
- [1332] T. Aaltonen (CDF), “Observation of Exclusive Dijet Production at the Fermilab Tevatron $p\bar{p}$ Collider”, Phys. Rev. D **77**, 052004 (2008).
- [1333] T. Aaltonen (CDF), “Observation of orbitally excited B_s mesons”, Phys. Rev. Lett. **100**, 082001 (2008).
- [1334] T. Aaltonen (CDF), “Observation of the Decay $B^+ \rightarrow J/\psi\pi^+$ and Measurement of the B^+ Mass”, Phys. Rev. Lett. **100**, 182002 (2008).
- [1335] T. Aaltonen (CDF), “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ decays with $2fb^{-1}$ of $p\bar{p}$ collisions”, Phys. Rev. Lett. **100**, 101802 (2008).
- [1336] T. Aaltonen (CDF), “Search for chargino-neutralino production in $p\bar{p}$ collisions at 1.96-TeV with high- $p(T)$ leptons”, Phys. Rev. D **77**, 052002 (2008).
- [1337] T. Aaltonen (CDF), “Search for Doubly Charged Higgs Bosons with Lepton-Flavor-Violating Decays involving Tau Leptons”, Phys. Rev. Lett. **101**, 121801 (2008).
- [1338] T. Aaltonen (CDF), “Search for Heavy Top-like Quarks Using Lepton Plus Jets Events in 1.96-TeV $p\bar{p}$ Collisions”, Phys. Rev. Lett. **100**, 161803 (2008).
- [1339] T. Aaltonen (CDF), “Search for Heavy, Long-Lived Neutralinos that Decay to Photons at CDF II Using Photon Timing”, Phys. Rev. D **78**, 032015 (2008).
- [1340] T. Aaltonen (CDF), “Search for large extra dimensions in final states containing one photon or jet and large missing transverse energy produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **101**, 181602 (2008).
- [1341] T. Aaltonen (CDF), “Search for New Heavy Particles Decaying to $Z^0Z^0 \rightarrow eeee$ in $p - \bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. D **78**, 012008 (2008).
- [1342] T. Aaltonen (CDF), “Search for Pair Production of Scalar Top Quarks Decaying to a τ Lepton and a b Quark in $p\bar{p}$ Collisions at $\sqrt{s}=1.96\text{ TeV}$ ”, Phys. Rev. Lett. **101**, 071802 (2008).
- [1343] T. Aaltonen (CDF), “Search for resonant $t\bar{t}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 231801 (2008).
- [1344] T. Aaltonen (CDF), “Search for Standard Model Higgs Boson Production in Association with a W Boson at CDF”, Phys. Rev. D **78**, 032008 (2008).
- [1345] T. Aaltonen (CDF), “Search for Standard Model Higgs Bosons Produced in Association with W Bosons”, Phys. Rev. Lett. **100**, 041801 (2008).
- [1346] T. Aaltonen (CDF), “Search for Supersymmetry in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ Using the Tripleton Signature of Chargino-Neutralino Production”, Phys. Rev. Lett. **101**, 251801 (2008).
- [1347] T. Aaltonen (CDF), “Search for the Flavor Changing Neutral Current Decay $t \rightarrow Zq$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **101**, 192002 (2008).
- [1348] T. Aaltonen (CDF), “Search for the Higgs boson in events with missing transverse energy and b quark jets produced in proton-antiproton collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **100**, 211801 (2008).
- [1349] T. Aaltonen (CDF), “Search for the Higgs boson produced with $Z \rightarrow \ell^+\ell^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **101**, 251803 (2008).

- [1350] T. Aaltonen (CDF), “Search for Third Generation Vector Leptoquarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **77**, 091105 (2008).
- [1351] T. Aaltonen (CDF), “Two-Particle Momentum Correlations in Jets Produced in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **77**, 092001 (2008).
- [1352] A. Abulencia (CDF), “Cross Section Measurements of High-p(T) Dilepton Final-State Processes Using a Global Fitting Method”, Phys. Rev. D **78**, 012003 (2008).
- [1353] A. Annovi, “A hardware track finder for the ATLAS trigger”, Nuovo Cim. B **123**, edited by M. Caffo, P. Capiluppi, U. Marconi, E. Remiddi, V. Vagnoni, and A. Zoccoli, 981–983 (2008).
- [1354] T. Aaltonen (CDF), “First measurement of the W boson mass in Run II of the Tevatron”, Phys. Rev. Lett. **99**, 151801 (2007).
- [1355] T. Aaltonen (CDF), “First observation of heavy baryons Σ_b and Σ_b^{*} ”, Phys. Rev. Lett. **99**, 202001 (2007).
- [1356] T. Aaltonen (CDF), “Limits on Anomalous Triple Gauge Couplings in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **76**, 111103 (2007).
- [1357] T. Aaltonen (CDF), “Measurement of the $p\bar{p} \rightarrow t\bar{t}$ production cross-section and the top quark mass at $\sqrt{s} = 1.96$ -TeV in the all-hadronic decay mode”, Phys. Rev. D **76**, 072009 (2007).
- [1358] T. Aaltonen (CDF), “Measurement of the top-quark mass in all-hadronic decays in p anti-p collisions at CDF II”, Phys. Rev. Lett. **98**, 142001 (2007).
- [1359] T. Aaltonen (CDF), “Measurement of the top-quark mass using missing E_T + jets events with secondary vertex b -tagging at CDF II”, Phys. Rev. D **75**, 111103 (2007).
- [1360] T. Aaltonen (CDF), “Model-Independent Global Search for New High-p(T) Physics at CDF”, 10.2172/922303 (2007).
- [1361] T. Aaltonen (CDF), “Observation and mass measurement of the baryon Xi_b^- ”, Phys. Rev. Lett. **99**, 052002 (2007).
- [1362] T. Aaltonen (CDF), “Search for a high-mass diphoton state and limits on Randall-Sundrum gravitons at CDF”, Phys. Rev. Lett. **99**, 171801 (2007).
- [1363] T. Aaltonen (CDF), “Search for chargino-neutralino production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **99**, 191806 (2007).
- [1364] T. Aaltonen (CDF), “Search for Direct Pair Production of Supersymmetric Top and Supersymmetric Bottom Quarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **76**, 072010 (2007).
- [1365] T. Aaltonen (CDF), “Search for exclusive $\gamma\gamma$ production in hadron-hadron collisions”, Phys. Rev. Lett. **99**, 242002 (2007).
- [1366] T. Aaltonen (CDF), “Search for New Particles Leading to Z + jets Final States in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **76**, 072006 (2007).
- [1367] T. Aaltonen (CDF), “Search for new physics in high mass electron-positron events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **99**, 171802 (2007).
- [1368] G. Aarons, “ILC Reference Design Report Volume 3 - Accelerator”, edited by N. Phinney, N. Toge, and N. J. Walker (2007).
- [1369] A. Abulencia (CDF), “Search for anomalous production of multi-lepton events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 131804 (2007).
- [1370] A. Abulencia (CDF), “Analysis of the quantum numbers J^{PC} of the $X(3872)$ ”, Phys. Rev. Lett. **98**, 132002 (2007).
- [1371] A. Abulencia (CDF), “First Measurement of the Ratio of Central-Electron to Forward-Electron W Partial Cross Sections in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **98**, 251801 (2007).
- [1372] A. Abulencia (CDF), “Inclusive Search for New Physics with Like-Sign Dilepton Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 221803 (2007).
- [1373] A. Abulencia (CDF), “Measurement of $\sigma_{\chi_{c2}}\mathcal{B}(\chi_{c2} \rightarrow J/\psi\gamma)/\sigma_{\chi_{c1}}\mathcal{B}(\chi_{c1} \rightarrow J/\psi\gamma)$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 232001 (2007).
- [1374] A. Abulencia (CDF), “Measurement of $\sigma(\Lambda_b^0)/\sigma(\bar{B}^0) \times \text{BR}(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)/\text{BR}(\bar{B}^0 \rightarrow D^+\pi^-)$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **98**, 122002 (2007).
- [1375] A. Abulencia (CDF), “Measurement of $\sigma(p\bar{p} \rightarrow Z) \cdot \text{Br}(Z \rightarrow 2\tau)$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **75**, 092004 (2007).
- [1376] A. Abulencia (CDF), “Measurement of the Λ_b^0 Lifetime in $\Lambda_b^0 \rightarrow J/\psi\Lambda^0$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 122001 (2007).
- [1377] A. Abulencia (CDF), “Measurement of the B^+ production cross-section in p anti-p collisions at $\sqrt{s} = 1.96$ -GeV”, Phys. Rev. D **75**, 012010 (2007).

- [1378] A. Abulencia (CDF), “Measurement of the Helicity Fractions of W Bosons from Top Quark Decays using Fully Reconstructed $t\bar{t}$ Events with CDF II”, Phys. Rev. D **75**, 052001 (2007).
- [1379] A. Abulencia (CDF), “Measurement of the Inclusive Jet Cross Section using the k_T algorithm in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV with the CDF II Detector”, Phys. Rev. D **75**, [Erratum: Phys.Rev.D 75, 119901 (2007)], 092006 (2007).
- [1380] A. Abulencia (CDF), “Measurement of the Ratios of Branching Fractions $B(B_0(s) \rightarrow D^{-(s)} \pi^+ \pi^+ \pi^-) / B(B_0 \rightarrow D^- \pi^+ \pi^+ \pi^-)$ and $B(B_0(s) \rightarrow D^{-(s)} \pi^+) / B(B_0 \rightarrow D^- \pi^+)$ ”, Phys. Rev. Lett. **98**, 061802 (2007).
- [1381] A. Abulencia (CDF), “Measurement of the Top Quark Mass in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV using the Decay Length Technique”, Phys. Rev. D **75**, 071102 (2007).
- [1382] A. Abulencia (CDF), “Measurements of inclusive W and Z cross sections in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV”, J. Phys. G **34**, 2457–2544 (2007).
- [1383] A. Abulencia (CDF), “Observation of WZ Production”, Phys. Rev. Lett. **98**, 161801 (2007).
- [1384] A. Abulencia (CDF), “Observation of Exclusive Electron-Positron Production in Hadron-Hadron Collisions”, Phys. Rev. Lett. **98**, 112001 (2007).
- [1385] A. Abulencia (CDF), “Polarization of J/ψ and ψ_{2S} Mesons Produced in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **99**, 132001 (2007).
- [1386] A. Abulencia (CDF), “Precise Measurement of the Top Quark Mass in the Lepton + Jets Topology at CDF II”, Phys. Rev. Lett. **99**, 182002 (2007).
- [1387] A. Abulencia (CDF), “Precision measurement of the top quark mass from dilepton events at CDF II”, Phys. Rev. D **75**, 031105 (2007).
- [1388] A. Abulencia (CDF), “Search for Exotic $S=-2$ Baryons in proton-antiproton Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **75**, 032003 (2007).
- [1389] A. Abulencia (CDF), “Search for Heavy, Long-Lived Particles that Decay to Photons at CDF II”, Phys. Rev. Lett. **99**, 121801 (2007).
- [1390] A. Abulencia (CDF), “Search for New Physics in Lepton + Photon + X Events with 929 pb^{-1} of $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **75**, 112001 (2007).
- [1391] A. Abulencia (CDF), “Search for $V+A$ current in top quark decay in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 072001 (2007).
- [1392] A. Abulencia (CDF), “Search for W-prime boson decaying to electron-neutrino pairs in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **75**, 091101 (2007).
- [1393] B. Aharmim (SNO), “Determination of the ν_e and total ^8B solar neutrino fluxes with the Sudbury neutrino observatory phase I data set”, Phys. Rev. C **75**, 045502 (2007).
- [1394] A. Abulencia (CDF), “A search for $t \rightarrow t\alpha\nu q$ in $t\bar{t}$ production.”, Phys. Lett. B **639**, 172–178 (2006).
- [1395] A. Abulencia (CDF), “A search for scalar bottom quarks from gluino decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 171802 (2006).
- [1396] A. Abulencia (CDF), “Direct search for Dirac magnetic monopoles in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **96**, 201801 (2006).
- [1397] A. Abulencia (CDF), “Evidence for the exclusive decay $B_c^\pm \rightarrow J/\psi\pi^\pm$ and measurement of the mass of the B_c meson”, Phys. Rev. Lett. **96**, 082002 (2006).
- [1398] A. Abulencia (CDF), “Measurement of mass and width of the excited charmed meson states $D_0(1)$ and $D_0^*(2)$ at CDF”, Phys. Rev. D **73**, 051104 (2006).
- [1399] A. Abulencia (CDF), “Measurement of the $B_s^0 - \bar{B}_s^0$ Oscillation Frequency”, Phys. Rev. Lett. **97**, 062003 (2006).
- [1400] A. Abulencia (CDF), “Measurement of the b jet cross-section in events with a Z boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. D **74**, 032008 (2006).
- [1401] A. Abulencia (CDF), “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **97**, 082004 (2006).
- [1402] A. Abulencia (CDF), “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV in the All Hadronic Decay Mode”, Phys. Rev. D **74**, 072005 (2006).
- [1403] A. Abulencia (CDF), “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV using Lepton + Jets Events with Jet Probability b^- tagging”, Phys. Rev. D **74**, 072006 (2006).
- [1404] A. Abulencia (CDF), “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV using missing E_T + jets events with secondary vertex b -tagging”, Phys. Rev. Lett. **96**, 202002 (2006).

- [1405] A. Abulencia (CDF), “Measurement of the $B(c)^+$ meson lifetime using $B(c)^+ \rightarrow J/\psi e^+ \nu(e)$ ”, Phys. Rev. Lett. **97**, 012002 (2006).
- [1406] A. Abulencia (CDF), “Measurement of the dipion mass spectrum in $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ decays.”, Phys. Rev. Lett. **96**, 102002 (2006).
- [1407] A. Abulencia (CDF), “Measurement of the helicity of W bosons in top-quark decays”, Phys. Rev. D **73**, 111103 (2006).
- [1408] A. Abulencia (CDF), “Measurement of the inclusive jet cross section in $p\bar{p}$ interactions at $\sqrt{s} = 1.96$ -TeV using a cone-based jet algorithm.”, Phys. Rev. D **74**, 071103 (2006).
- [1409] A. Abulencia (CDF), “Measurement of the inclusive jet cross section using the $k(t)$ algorithm in p anti- p collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 122001 (2006).
- [1410] A. Abulencia (CDF), “Measurement of the ratio of branching fractions $B(D0 \rightarrow K^+ \pi^-) / B(D0 \rightarrow K^- \pi^+)$ using the CDF II Detector”, Phys. Rev. D **74**, 031109 (2006).
- [1411] A. Abulencia (CDF), “Measurement of the ratios of branching fractions $B(B_s^0 \rightarrow D_s^- \pi^+) / B(B^0 \rightarrow D^- \pi^+)$ and $B(B^+ \rightarrow \bar{D}^0 \pi^+) / B(B^0 \rightarrow D^- \pi^+)$ ”, Phys. Rev. Lett. **96**, 191801 (2006).
- [1412] A. Abulencia (CDF), “Measurement of the top quark mass using template methods on dilepton events in proton antiproton collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. D **73**, 112006 (2006).
- [1413] A. Abulencia (CDF), “Measurement of the top quark mass with the dynamical likelihood method using lepton plus jets events with b-tags in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. D **73**, 092002 (2006).
- [1414] A. Abulencia (CDF), “Observation of $B_s^0 \rightarrow \psi(2S)\phi$ and measurement of ratio of branching fractions $B(B_s^0 \rightarrow \psi(2S)\phi) / B(B_s^0 \rightarrow J/\psi\phi)$ ”, Phys. Rev. Lett. **96**, 231801 (2006).
- [1415] A. Abulencia (CDF), “Observation of $B_s^0 - \bar{B}_s^0$ Oscillations”, Phys. Rev. Lett. **97**, 242003 (2006).
- [1416] A. Abulencia (CDF), “Observation of $B^0 (s) \rightarrow K^+ K^-$ and Measurements of Branching Fractions of Charmless Two-body Decays of B^0 and B_s^0 Mesons in $\bar{p}p$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 211802 (2006).
- [1417] A. Abulencia (CDF), “Precision top quark mass measurement in the lepton + jets topology in p anti- p collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 022004 (2006).
- [1418] A. Abulencia (CDF), “Search for $Z' \rightarrow e^+ e^-$ using dielectron mass and angular distribution.”, Phys. Rev. Lett. **96**, 211801 (2006).
- [1419] A. Abulencia (CDF), “Search for a neutral Higgs boson decaying to a W boson pair in p anti- p collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 081802 (2006).
- [1420] A. Abulencia (CDF), “Search for anomalous semileptonic decay of heavy flavor hadrons produced in association with a W boson at CDF II”, Phys. Rev. D **73**, 051101 (2006).
- [1421] A. Abulencia (CDF), “Search for charged Higgs bosons from top quark decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 042003 (2006).
- [1422] A. Abulencia (CDF), “Search for excited and exotic muons in the $\mu\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 191802 (2006).
- [1423] A. Abulencia (CDF), “Search for H to b anti- b produced in association with W bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 081803 (2006).
- [1424] A. Abulencia (CDF), “Search for high-mass resonances decaying to $e \mu$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 211802 (2006).
- [1425] A. Abulencia (CDF), “Search for Large Extra Dimensions in the Production of Jets and Missing Transverse Energy in p anti- p Collisions at $s^{*(1/2)} = 1.96$ TeV”, Phys. Rev. Lett. **97**, 171802 (2006).
- [1426] A. Abulencia (CDF), “Search for neutral MSSM Higgs bosons decaying to tau pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **96**, 011802 (2006).
- [1427] A. Abulencia (CDF), “Search for new physics in lepton + photon + X events with 305 pb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 031801 (2006).
- [1428] A. Abulencia (CDF), “Search for second-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. D **73**, 051102 (2006).
- [1429] A. Abulencia (CDF), “Top quark mass measurement from dilepton events at CDF II”, Phys. Rev. Lett. **96**, 152002 (2006).
- [1430] A. Abulencia (CDF), “Top quark mass measurement from dilepton events at CDF II with the matrix-element method”, Phys. Rev. D **74**, 032009 (2006).
- [1431] A. Abulencia (CDF), “Top quark mass measurement using the template method in the lepton + jets channel at CDF II”, Phys. Rev. D **73**, 032003 (2006).

- [1432] D. Acosta (CDF), “Measurement of b hadron masses in exclusive J/ψ decays with the CDF detector”, Phys. Rev. Lett. **96**, 202001 (2006).
- [1433] D. Acosta (CDF), “Measurement of the azimuthal angle distribution of leptons from W boson decays as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **73**, 052002 (2006).
- [1434] D. Acosta (CDF), “Search for W and Z bosons in the reaction $\bar{p}p \rightarrow 2\text{jets} + \gamma$ at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **73**, 012001 (2006).
- [1435] A. Abulencia (CDF), “Search for $B_s \rightarrow \mu^+\mu^-$ and $B_d \rightarrow \mu^+\mu^-$ decays in $p\bar{p}$ collisions with CDF II”, Phys. Rev. Lett. **95**, [Erratum: Phys.Rev.Lett. 95, 249905 (2005)], 221805 (2005).
- [1436] A. Abulencia (CDF), “Search for new high mass particles decaying to lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **95**, 252001 (2005).
- [1437] D. Acosta (CDF), “ K_S^0 and Λ^0 production studies in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ -GeV and 630 -GeV”, Phys. Rev. D **72**, 052001 (2005).
- [1438] D. Acosta (CDF), “A search for supersymmetric Higgs bosons in the di-tau decay mode in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **72**, 072004 (2005).
- [1439] D. Acosta (CDF), “Comparison of three-jet events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV to predictions from a next-to-leading order QCD calculation”, Phys. Rev. D **71**, 032002 (2005).
- [1440] D. Acosta (CDF), “First evidence for $B_s^0 \rightarrow \phi\phi$ decay and measurements of branching ratio and A_{CP} for $B^+ \rightarrow \phi K^+$ ”, Phys. Rev. Lett. **95**, 031801 (2005).
- [1441] D. Acosta (CDF), “First measurements of inclusive W and Z cross sections from Run II of the Tevatron collider”, Phys. Rev. Lett. **94**, 091803 (2005).
- [1442] D. Acosta (CDF), “Measurement of $B(t \rightarrow Wb)/B(t \rightarrow Wq)$ at the Collider Detector at Fermilab”, Phys. Rev. Lett. **95**, 102002 (2005).
- [1443] D. Acosta (CDF), “Measurement of $W\gamma$ and $Z\gamma$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **94**, 041803 (2005).
- [1444] D. Acosta (CDF), “Measurement of charged particle multiplicities in gluon and quark jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **94**, 171802 (2005).
- [1445] D. Acosta (CDF), “Measurement of partial widths and search for direct CP violation in D^0 meson decays to K^-K^+ and $\pi^-\pi^+$ ”, Phys. Rev. Lett. **94**, 122001 (2005).
- [1446] D. Acosta (CDF), “Measurement of the J/ψ meson and b -hadron production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1960$ GeV”, Phys. Rev. D **71**, 032001 (2005).
- [1447] D. Acosta (CDF), “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using kinematic fitting of b -tagged lepton + jet events”, Phys. Rev. D **71**, 072005 (2005).
- [1448] D. Acosta (CDF), “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using lepton + jets events with secondary vertex b -tagging”, Phys. Rev. D **71**, 052003 (2005).
- [1449] D. Acosta (CDF), “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using lepton plus jets events with semileptonic B decays to muons”, Phys. Rev. D **72**, 032002 (2005).
- [1450] D. Acosta (CDF), “Measurement of the W^+W^- production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using dilepton events”, Phys. Rev. Lett. **94**, 211801 (2005).
- [1451] D. Acosta (CDF), “Measurement of the W boson polarization in top decay at CDF at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **71**, [Erratum: Phys.Rev.D 71, 059901 (2005)], 031101 (2005).
- [1452] D. Acosta (CDF), “Measurement of the cross section for $t\bar{t}$ production in $p\bar{p}$ collisions using the kinematics of lepton + jets events”, Phys. Rev. D **72**, 052003 (2005).
- [1453] D. Acosta (CDF), “Measurement of the cross section for prompt diphoton production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **95**, 022003 (2005).
- [1454] D. Acosta (CDF), “Measurement of the forward-backward charge asymmetry from $W \rightarrow e\nu$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **71**, 051104 (2005).
- [1455] D. Acosta (CDF), “Measurement of the forward-backward charge asymmetry of electron positron pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **71**, 052002 (2005).
- [1456] D. Acosta (CDF), “Measurement of the lifetime difference between B_s mass eigenstates”, Phys. Rev. Lett. **94**, 101803 (2005).
- [1457] D. Acosta (CDF), “Measurement of the moments of the hadronic invariant mass distribution in semileptonic B decays”, Phys. Rev. D **71**, 051103 (2005).
- [1458] D. Acosta (CDF), “Measurements of $b\bar{b}$ azimuthal production correlations in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **71**, 092001 (2005).

- [1459] D. Acosta (CDF), “Search for $\Lambda_b \rightarrow p\pi$ and $\Lambda_b \rightarrow pK$ decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **72**, 051104 (2005).
- [1460] D. Acosta (CDF), “Search for ZZ and ZW production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **71**, 091105 (2005).
- [1461] D. Acosta (CDF), “Search for anomalous kinematics in $t\bar{t}$ dilepton events at CDF II”, Phys. Rev. Lett. **95**, 022001 (2005).
- [1462] D. Acosta (CDF), “Search for anomalous production of diphoton events with missing transverse energy at CDF and limits on gauge-mediated supersymmetry-breaking models”, Phys. Rev. D **71**, 031104 (2005).
- [1463] D. Acosta (CDF), “Search for electroweak single top quark production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **71**, 012005 (2005).
- [1464] D. Acosta (CDF), “Search for excited and exotic electrons in the $e\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **94**, 101802 (2005).
- [1465] D. Acosta (CDF), “Search for first-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **72**, 051107 (2005).
- [1466] D. Acosta (CDF), “Search for Higgs bosons decaying into $b\bar{b}$ and produced in association with a vector boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **95**, 051801 (2005).
- [1467] D. Acosta (CDF), “Search for long-lived doubly-charged Higgs bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **95**, 071801 (2005).
- [1468] D. Acosta (CDF), “Search for new physics using high mass tau pairs from 1.96 TeV $p\bar{p}$ collisions”, Phys. Rev. Lett. **95**, 131801 (2005).
- [1469] D. Acosta (CDF), “Search for scalar leptoquark pairs decaying to $\nu\bar{\nu}q\bar{q}$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **71**, [Erratum: Phys.Rev.D 71, 119901 (2005)], 112001 (2005).
- [1470] D. Acosta (CDF), “Study of jet shapes in inclusive jet production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **71**, 112002 (2005).
- [1471] B. Aharmim (SNO), “Electron energy spectra, fluxes, and day-night asymmetries of B-8 solar neutrinos from measurements with NaCl dissolved in the heavy-water detector at the Sudbury Neutrino Observatory”, Phys. Rev. C **72**, 055502 (2005).
- [1472] M. Casarsa, S. C. Hsu, E. D. Lipeles, **M. Neubauer**, S. Sarkar, I. Sfiligoi, F. K. Wuerthwein, F. Wuerthwein, “The CDF analysis farm”, AIP Conf. Proc. **794**, edited by A. Tricomi, S. Albergo, and M. Chiorboli, 275–278 (2005).
- [1473] V. M. Abazov (CDF, D0), “Combination of CDF and D0 Results on W Boson Mass and Width”, Phys. Rev. D **70**, 092008 (2004).
- [1474] D. Acosta (CDF), “Direct photon cross section with conversions at CDF”, Phys. Rev. D **70**, 074008 (2004).
- [1475] D. Acosta (CDF), “Inclusive double pomeron exchange at the Fermilab Tevatron $p\bar{p}$ collider”, Phys. Rev. Lett. **93**, 141601 (2004).
- [1476] D. Acosta (CDF), “Inclusive search for anomalous production of high p_T like-sign lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **93**, 061802 (2004).
- [1477] D. Acosta (CDF), “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using dilepton events”, Phys. Rev. Lett. **93**, 142001 (2004).
- [1478] D. Acosta (CDF), “Measurement of the average time-integrated mixing probability of b -flavored hadrons produced at the Tevatron”, Phys. Rev. D **69**, 012002 (2004).
- [1479] D. Acosta (CDF), “Measurement of the polar-angle distribution of leptons from W boson decay as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **70**, 032004 (2004).
- [1480] D. Acosta (CDF), “Observation of the narrow state $X(3872) \rightarrow J/\psi\pi^+\pi^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **93**, 072001 (2004).
- [1481] D. Acosta (CDF), “Optimized search for single top quark production at the Fermilab Tevatron”, Phys. Rev. D **69**, 052003 (2004).
- [1482] D. Acosta (CDF), “Search for doubly-charged Higgs bosons decaying to dileptons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **93**, 221802 (2004).
- [1483] D. Acosta (CDF), “Search for Kaluza-Klein graviton emission in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV using the missing energy signature”, Phys. Rev. Lett. **92**, 121802 (2004).
- [1484] D. Acosta (CDF), “Search for pair production of scalar top quarks in R -parity violating decay modes in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **92**, 051803 (2004).

- [1485] D. Acosta (CDF), “The underlying event in hard interactions at the Tevatron $p\bar{p}$ collider”, Phys. Rev. D **70**, 072002 (2004).
- [1486] D. Acosta (CDF), “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **93**, 032001 (2004).
- [1487] B. Aharmim (SNO), “Electron antineutrino search at the Sudbury Neutrino Observatory”, Phys. Rev. D **70**, 093014 (2004).
- [1488] S. N. Ahmed (SNO), “Constraints on nucleon decay via ‘invisible’ modes from the Sudbury Neutrino Observatory”, Phys. Rev. Lett. **92**, 102004 (2004).
- [1489] T. H. Kim, **M. Neubauer**, I. Sfiligoi, L. Weems, F. K. Wuerthwein, “The CDF Central Analysis Farm”, IEEE Trans. Nucl. Sci. **51**, edited by J. P. Dufey, 892–896 (2004).
- [1490] D. Acosta (CDF), “Central pseudorapidity gaps in events with a leading antiproton at the Fermilab Tevatron $p\bar{p}$ collider”, Phys. Rev. Lett. **91**, 011802 (2003).
- [1491] D. Acosta (CDF), “Measurement of prompt charm meson production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **91**, 241804 (2003).
- [1492] D. Acosta (CDF), “Measurement of the mass difference $m(D_s^+) - m(D^+)$ at CDF II”, Phys. Rev. D **68**, 072004 (2003).
- [1493] D. Acosta (CDF), “Search for a W' Boson Decaying to a Top and Bottom Quark Pair in 1.8 TeV $p\bar{p}$ Collisions”, Phys. Rev. Lett. **90**, 081802 (2003).
- [1494] D. Acosta (CDF), “Search for Associated Production of Υ and Vector Boson in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **90**, 221803 (2003).
- [1495] D. Acosta (CDF), “Search for lepton flavor violating decays of a heavy neutral particle in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **91**, 171602 (2003).
- [1496] D. Acosta (CDF), “Search for Long-Lived Charged Massive Particles in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **90**, 131801 (2003).
- [1497] D. Acosta (CDF), “Search for the flavor-changing neutral current decay $D^0 \rightarrow \mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. D **68**, 091101 (2003).
- [1498] D. Acosta (CDF), “Search for the supersymmetric partner of the top quark in dilepton events from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **90**, 251801 (2003).
- [1499] **M. S. Neubauer** (CDF), “Computing for Run II at CDF”, Nucl. Instrum. Meth. A **502**, edited by V. A. Ilyin, V. V. Korenkov, and D. Perret-Gallix, 386–390 (2003).
- [1500] D. Acosta (CDF), “Cross Section for Forward J/ψ Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. D **66**, 092001 (2002).
- [1501] Q. R. Ahmad (SNO), “Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory”, Phys. Rev. Lett. **89**, 011301 (2002).
- [1502] Q. R. Ahmad (SNO), “Measurement of day and night neutrino energy spectra at SNO and constraints on neutrino mixing parameters”, Phys. Rev. Lett. **89**, 011302 (2002).
- [1503] A. B. McDonald (SNO), “Direct evidence for neutrino flavor transformation from neutral-current interactions in SNO”, AIP Conf. Proc. **646**, edited by V. Elias, R. J. Epp, and R. C. Myers, 43–58 (2002).
- [1504] A. W. P. Poon (SNO), “Neutrino observations from the Sudbury Neutrino Observatory”, AIP Conf. Proc. **610**, edited by E. Norman, L. Schroeder, and G. Wozniak, 218–230 (2002).
- [1505] A. W. P. Poon (SNO), “Solar neutrino observations at the Sudbury Neutrino Observatory”, eConf **C020805**, edited by J. Hewett, J. Jaros, T. Kamae, and C. Prescott, TTH01 (2002).
- [1506] Q. R. Ahmad (SNO), “Measurement of the rate of $\nu_e + d \rightarrow p + p + e^-$ interactions produced by ^8B solar neutrinos at the Sudbury Neutrino Observatory”, Phys. Rev. Lett. **87**, 071301 (2001).
- [1507] A. B. McDonald (SNO), “Measurement of CC interactions produced by ^8B solar neutrinos at SNO”, PoS **hep2001**, edited by D. Horváth, P. Lévai, and A. Patkós, 186 (2001).
- [1508] **M. S. Neubauer** (SNO), “First solar neutrino observations from SNO”, Int. J. Mod. Phys. A **16S1B**, edited by K. K. Gan and R. Kass, 715–717 (2001).
- [1509] J. Boger (SNO), “The Sudbury neutrino observatory”, Nucl. Instrum. Meth. A **449**, 172–207 (2000).

Conference Proceedings

- [1510] G. Benelli, “Data Science and Machine Learning in Education”, 2022 Snowmass Summer Study (July 2022).

- [1511] P. Harris, “Physics Community Needs, Tools, and Resources for Machine Learning”, 2022 Snowmass Summer Study (Mar. 2022).
- [1512] A. Lazar, “Accelerating the Inference of the Exa.TrkX Pipeline”, 20th International Workshop on Advanced Computing and Analysis Techniques in Physics Research: AI Decoded - Towards Sustainable, Diverse, Performant and Effective Scientific Computing (Feb. 2022).
- [1513] **M. S. Neubauer**, A. Roy, “Explainable AI for High Energy Physics”, 2022 Snowmass Summer Study (June 2022).
- [1514] S. Thais, P. Calafiura, G. Chachamis, G. DeZoort, J. Duarte, S. Ganguly, M. Kagan, D. Murnane, **M. S. Neubauer**, K. Terao, “Graph Neural Networks in Particle Physics: Implementations, Innovations, and Challenges”, 2022 Snowmass Summer Study (Mar. 2022).
- [1515] C.-Y. Wang, “Reconstruction of Large Radius Tracks with the Exa.TrkX pipeline”, 20th International Workshop on Advanced Computing and Analysis Techniques in Physics Research: AI Decoded - Towards Sustainable, Diverse, Performant and Effective Scientific Computing (Mar. 2022).
- [1516] **M. S. Neubauer**, “Learning from the Pandemic: the Future of Meetings in HEP and Beyond”, IRIS-HEP Blueprint Workshop (June 2021).
- [1517] T. Aarrestad (HEP Software Foundation), “HL-LHC Computing Review: Common Tools and Community Software”, 2022 Snowmass Summer Study, edited by P. Canal et al. (Aug. 2020).
- [1518] A. Heintz, “Accelerated Charged Particle Tracking with Graph Neural Networks on FPGAs”, 34th Conference on Neural Information Processing Systems (Nov. 2020).
- [1519] G. Allen, “Deep Learning for Multi-Messenger Astrophysics: A Gateway for Discovery in the Big Data Era”, (Feb. 2019).
- [1520] G. Kazakevich, R. Johnson, **M. Neubauer**, V. Lebedev, W. Schappert, V. Yakovlev, “Methods of phase and power control in magnetron transmitters for superconducting accelerators”, 8th International Particle Accelerator Conference (May 2017).
- [1521] T. Aaltonen (CDF, D0), “Combined CDF and D0 Upper Limits on Standard Model Higgs Boson Production with up to 8.2 fb^{-1} of Data”, 46th Rencontres de Moriond on Electroweak Interactions and Unified Theories (Mar. 2011).
- [1522] **M. S. Neubauer** (ATLAS), “A Fast Hardware Tracker for the ATLAS Trigger System”, Meeting of the APS Division of Particles and Fields (Oct. 2011).
- [1523] A. Annovi, “The Evolution of FTK, a Real-Time Tracker for Hadron Collider Experiments”, 11th ICATPP Conference on Astroparticle, Particle, Space Physics, Detectors and Medical Physics Applications (2010), pp. 653–657.
- [1524] **M. S. Neubauer** (ATLAS TDAQ), “A fast hardware tracker for the ATLAS trigger system”, 2009 IEEE Nuclear Science Symposium and Medical Imaging Conference (2009), pp. 1876–1879.
- [1525] **M. S. Neubauer** (CDF, D0), “Diboson physics at the Tevatron”, 41st Rencontres de Moriond: QCD and Hadronic Interactions (May 2006), pp. 113–116.
- [1526] A. Sill, “Globally distributed user analysis computing at CDF”, 14th International Conference on Computing in High-Energy and Nuclear Physics (2005), pp. 1139–1142.
- [1527] F. K. Wuerthwein, E. D. Lipeles, **M. Neubauer**, I. Sfiligoi, “The Condor based CDF CAF”, 14th International Conference on Computing in High-Energy and Nuclear Physics (2005), pp. 978–981.
- [1528] A. McDonald, “First results from the Sudbury Neutrino Observatory”, NO-VE International Workshop on Neutrino Oscillations in Venice (July 2001), pp. 1–16.